

# The statistics of triggered star formation: an overdensity of massive young stellar objects around *Spitzer* bubbles

M. A. Thompson,<sup>1</sup>★ J. S. Urquhart,<sup>2,3</sup> T. J. T. Moore<sup>4</sup> and L. K. Morgan<sup>4</sup>

<sup>1</sup>Centre for Astrophysics Research, Science and Technology Research Institute, University of Hertfordshire, College Lane, Hatfield, Herts AL10 9AB

<sup>2</sup>Australia Telescope National Facility, CSIRO Astronomy and Space Science, PO Box 76, Epping NSW 1710, Australia

<sup>3</sup>Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

<sup>4</sup>Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead CH41 1LD

Accepted 2011 December 1. Received 2011 December 1; in original form 2011 November 3

## ABSTRACT

We present a detailed statistical study of massive star formation in the environment of 322 *Spitzer* mid-infrared bubbles by using the Red MSX Source (RMS) survey for massive young stellar objects (YSOs). Using a combination of simple surface density plots and a more sophisticated angular cross-correlation function analysis, we show that there is a statistically significant overdensity of RMS YSOs towards the bubbles. There is a clear peak in the surface density and angular cross-correlation function of YSOs projected against the rim of the bubbles. By investigating the autocorrelation function of the RMS YSOs, we show that this is not due to intrinsic clustering of the RMS YSO sample. RMS YSOs and *Spitzer* bubbles are essentially uncorrelated with each other beyond a normalized angular distance of two bubble radii. The bubbles associated with RMS YSOs tend to be both smaller and thinner than those that are not associated with YSOs. We interpret this tendency to be due to an age effect, with YSOs being preferentially found around smaller and younger bubbles. We find no evidence to suggest that the YSOs associated with the bubbles are any more luminous than the rest of the RMS YSO population, which suggests that the triggering process does not produce a top-heavy luminosity function or initial mass function. We suggest that it is likely that the YSOs were triggered by the expansion of the bubbles and estimate that the fraction of massive stars in the Milky Way formed by this process could be between 14 and 30 per cent.

**Key words:** stars: formation – ISM: bubbles – H II regions – infrared: ISM.

## 1 INTRODUCTION

Massive stars drive large amounts of energy into the surrounding interstellar medium (ISM) via their winds and ionizing radiation, sculpting the ISM into a series of bubble and shell-like structures (Elmegreen 2011). One particular class of shells or bubbles are those surrounding expanding H II regions. The ultraviolet (UV) illumination excites emission from polycyclic aromatic hydrocarbons (PAHs) at the bubble rims and also heats dust grains within the H II region. The former can be readily observed at 8  $\mu$ m by IRAC on-board the *Spitzer Space Telescope* (Churchwell et al. 2006, 2007) and the latter at longer wavelengths using e.g. MIPS on *Spitzer* (Churchwell et al. 2007; Watson et al. 2008) or PACS on-board the *Herschel Space Observatory* (Zavagno et al. 2010a,b).

The expansion of these H II regions is of extreme interest to studies of star formation as their expansion may trigger new generations of star formation into being within the molecular material surrounding

the bubbles. There are two major triggering mechanisms that have been put forward so far: radiative-driven implosion (RDI) and the collect and collapse process. In RDI (Bertoldi 1989; Lefloch & Lazareff 1994; Bisbas et al. 2009, 2011; Miao et al. 2009) the expanding ionization front of the H II region drives a D-type shock into molecular clouds surrounding the H II region, triggering the collapse of subcritical clumps within the clouds. Theoretical models of RDI can successfully explain the morphology of bright-rimmed clouds (BRCs) (Bisbas et al. 2009, 2011; Miao et al. 2009) and observations suggest that star formation is concentrated along the central axis of the clouds (Sugitani, Tamura & Ogura 1999; Sugitani et al. 2000; Bisbas et al. 2011).

The collect and collapse process (Elmegreen & Lada 1977; Whitworth et al. 1994; Whitworth & Francis 2002), on the other hand, does not require the presence of pre-existing molecular structures. In this case, the expansion of the H II region sweeps up the surrounding low-density material into a shell surrounding the H II region. At a certain point (generally after a few Myr), this shell becomes self-gravitating, fragments and collapses to form dense molecular clumps that eventually collapse to form stars. The

★E-mail: m.a.thompson@herts.ac.uk


fragments formed in the collect and collapse process tend to be massive (a few hundred  $M_{\odot}$ ; Whitworth et al. 1994) and so this process could naturally explain the hierarchical nature of massive stellar clusters (e.g. Bastian et al. 2005; Oey et al. 2005). It should be noted that neither mechanism excludes the other as a means of triggering star formation. Indeed, within a single H II region, both RDI and collect and collapse may operate together (Deharveng, Zavagno & Caplan 2005).

However, recent models of the ionizing feedback from massive stars (Dale & Bonnell 2011) suggest that neither mechanism is responsible and that the UV illumination from these stars simply erodes low-density material rather than shaping its evolution and eventual collapse. In this scenario, the role of triggering in star formation is predicted to be minimal. Walch et al. (2011) find that the location of the dense clumps around the edge of H II regions (such as RCW 120) reflects the pre-existing cloud structure and their formation does not require the collect and collapse process. However, rather than neglecting triggering as in Dale & Bonnell (2011), they suggest that stars may form by global implosion of the pre-existing structures (Enhancement of initial Density substructure and simultaneous Global Implosion or EDGI).

Observational studies of triggered star formation have so far mainly focused on photoionized globules (BRCs; Sugitani, Fukui & Ogura 1991; Sugitani & Ogura 1994) found at the edges of optically visible H II regions (Morgan et al. 2004, 2008; Thompson et al. 2004; Urquhart et al. 2004, 2006) or on the rims of H II regions selected to show a relatively simple morphology (Deharveng et al. 2005; Zavagno et al. 2006; Pomarès et al. 2009). Recent studies of the latter have been able to take advantage of the large catalogue of infrared bubbles discovered in the GLIMPSE survey (Churchwell et al. 2006, 2007). The properties of these bubbles are consistent with expanding H II regions (Watson et al. 2009; Deharveng et al. 2010), although perhaps more consistent with a ring rather than bubble morphology (Beaumont & Williams 2010).

However, one difficulty with many of these studies is that they are often phenomenological in nature, concentrating upon the visual identification of YSOs or protostars in regions where one might have a prior expectation that they may have been triggered (e.g. Zavagno et al. 2006, 2007; Deharveng et al. 2010). Such studies cannot attack the central problem in triggered star formation, which is to identify the origin of the discovered star formation (we refer to this as the “origin problem”). When trying to identify star formation as being triggered, one must first exclude the possibility that the stars would have formed spontaneously without the influence of the trigger. The origin problem is particularly intractable when considering individual objects – without a good understanding of the initial conditions involved it is almost impossible to categorize an individual star-forming region as being triggered or spontaneous.

In this paper, we attempt to move beyond the current phenomenological approach by carrying out a detailed statistical study of star formation around the 322 *Spitzer* bubbles of Churchwell et al. (2006). While the origin problem is almost impossible to solve for individual star-forming regions, by considering the global properties of a large sample it may be possible to infer the presence of triggering in a statistical sense. These studies are made easier by the existence of large, uniformly selected and well-understood star formation surveys such as the Red MSX Source (RMS) survey (Urquhart et al. 2008). The RMS survey has the goal of identifying every massive young stellar object (YSO) in the Milky Way and comprises an initial infrared selection followed by a thorough multiwavelength follow-up to rigorously classify each object and



**Figure 1.** Three-colour GLIMPSE image of a *Spitzer* bubble (N109 from the Churchwell et al. 2006 catalogue), showing the strong extended 8- $\mu$ m PAH and 24- $\mu$ m dust emission tracing the bubble rim. The colour coding in the image is 24  $\mu$ m (red), 8  $\mu$ m (green) and 4.5  $\mu$ m (blue). The 24- $\mu$ m image is taken from the MIPS GAL survey (Carey et al. 2009). N109 is one of the largest bubbles in the Churchwell et al. (2006) catalogue, with a mean radius of 14.8 arcmin, and is also the site of numerous smaller bubbles. The positions of two objects from the RMS YSO sample are indicated by circles.

determine its physical properties (Urquhart et al. 2009b,c, 2011; Mottram et al. 2011a).

The RMS survey covers a greater area than the GLIMPSE-I survey region in which the Churchwell et al. (2006) bubbles have been identified and is complete to YSOs of luminosity  $\geq 10^4 L_{\odot}$  out to the furthest bubble in the Churchwell et al. (2006) catalogue. Hence, we can conduct a survey for recent massive star formation around *all* of the *Spitzer* bubbles (see Fig. 1 for an example). Note that the RMS survey does not include Galactic latitudes  $|l| \leq 10^\circ$  for reasons of confusion and so we do not consider here the bubble catalogue of Churchwell et al. (2007), which is solely based upon the GLIMPSE-II survey region (i.e.  $|l| \leq 10^\circ$ ). On the other hand, we must keep in mind the intrinsic subjective biases in the catalogue of *Spitzer* bubbles, due to their manual by-eye identification by a number of independent observers (Churchwell et al. 2006). These biases are discussed in detail in Section 2 of Churchwell et al. (2006). The current catalogue is likely to be highly incomplete, particularly to small bubbles. Churchwell et al. (2006) estimate a completeness of the order of  $\sim 50$  per cent, which is being borne out by early results from the citizen science Milky Way Project<sup>1</sup> (Simpson et al., in preparation).

Nevertheless, the Churchwell et al. (2006) catalogue of bubbles currently represents the most complete and well-studied catalogue of H II regions with a simple morphology that lends well to statistical studies of their YSO distribution. In this paper, we present such a study, paying particular attention to the angular distribution of YSOs around the bubbles and potential differences within the YSO population. Our aims are to provide statistical evidence that the star formation associated with the bubbles may have been triggered and to investigate the properties of the global population of bubbles and YSOs. In addition, the methods that we demonstrate in this paper will be readily applicable to the larger and more complete Milky Way Project bubble sample when it becomes available.

<sup>1</sup> <http://www.milkywayproject.org>

This paper is structured as follows. In Section 2 we study the star-forming environment of the *Spitzer* bubbles using a simple surface density approach, followed by a more sophisticated analysis of the angular cross-correlation function of RMS YSOs and *Spitzer* bubbles. Section 3 combines the results of the star-forming environment analysis with the observed properties of the bubbles and YSOs to demonstrate that star formation is clearly enhanced towards the bubbles. We speculate on the likely origin of this star formation and show that it is likely that the bubbles predate the formation of the RMS YSOs. Finally, in Section 4, we present a summary of our conclusions and results.

## 2 THE STAR-FORMING ENVIRONMENT OF *SPITZER* BUBBLES


### 2.1 The surface density of YSOs

As a first approach to studying the distribution of RMS YSOs around *Spitzer* bubbles, we simply measured the number counts of RMS YSOs expressed as a function of angular separation from the bubble centres. Our sample of RMS YSOs is comprised of the objects classified as either YSO or ultracompact (UC) H II in the RMS data base<sup>2</sup> (see Urquhart et al. 2008 for a description of the RMS data base and its classification system). YSO and UC H II sub-classifications both represent young, recently formed and predominantly massive stellar objects that enable us to trace the distribution of recent star formation around the bubbles. Hereafter we refer to this combined population as the RMS YSO sample. Within the area covered by the GLIMPSE-I survey (Benjamin et al. 2003), there are 846 objects within the RMS YSO sample and 322 bubbles from the Churchwell et al. (2006) catalogue.

In order to account for the different angular radii of the bubbles, we divided the angular separation of each RMS object from a particular bubble by the mean radius of the bubble ( $\langle R \rangle$ , column 9 in the catalogue of Churchwell et al. 2006), i.e. expressing the angular separation in bubble radii rather than arcminutes. Each bin represents an annulus around the centre of each bubble. The surface area of the annulus thus naturally increases with increasing radius and so to obtain the surface density of RMS YSOs we scale the counts in each bin by the surface area of each corresponding annulus. A histogram of these scaled number counts is shown in Fig. 2.

Fig. 2 shows a clear peak in the number of RMS YSOs at a separation of 1 normalized bubble radius. At greater angular radii, the number of RMS YSOs falls sharply, reaching a constant background level of  $\sim 3$  sources by 2 bubble radii. Within an angular radius of 2 normalized bubble radii, the number of RMS YSOs is demonstrably higher than at that angular radii greater than 2 bubble radii. The surface density of YSOs projected against *Spitzer* bubbles is thus higher than regions external to the bubbles, with a clear peak in the surface density projected against the rims of the bubbles.

The *Spitzer* bubbles are relatively elliptical, with typical eccentricities between 0.6 and 0.7. As we normalize by the mean bubble radius ( $\langle R \rangle$ ), this will have the effect that we incorrectly calculate the true normalized radius of each RMS YSO from the bubble centre, potentially broadening the observed peak in surface density. The position angles of the elliptical fits to the bubbles are not listed in Churchwell et al. (2006), but these measurements were kindly made available by Matt Povich (private communication) so that we could examine the effect of using the true radius of the bubble instead



**Figure 2.** Histogram of the number counts of RMS YSOs (comprising YSO and UC H II classifications from the RMS data base) as a function of angular distance from the centre of *Spitzer* bubbles. The distance is expressed in terms of the normalized bubble radius. The number counts are scaled by the area of the annulus corresponding to each bin and thus represent a surface density of RMS YSOs. Error bars are determined via Poisson statistics.


of the mean radius. We found that there is no significant difference between scaling the distance of the RMS YSOs with the true bubble radius and the mean radius. This is more than likely due to the fact that the angular resolution of our histogram in Fig. 2 is limited to 0.25 bubble radii by the need to obtain sufficient RMS YSOs in each bin. At this resolution, the worst-case error in radius (i.e. between mean radius  $\langle R \rangle$  and the semimajor axis  $a$ ) is slightly larger than the width of one bin in Fig. 2.

We further subdivided our RMS YSO sample into its constituent YSO and UC H II subsamples to investigate trends in the separate distributions of YSOs and UC H IIs, for example in evolutionary status versus radius. We found that there is no significant difference between the two subsamples, the histograms of separate YSO and UC H II subsamples are indistinguishable from the combined RMS YSO sample. Again, this may be due to the limited sample statistics that we currently have, or this may indicate that gradients in evolutionary status around the bubbles are either not present or, if present, are on smaller angular scales than resolved by our study.

In order to confirm this result, we also carried out the same analysis on the Robitaille et al. (2008) catalogue of intrinsically red sources selected from the *Spitzer* GLIMPSE survey. The Robitaille catalogue contains a much greater number of objects than the RMS survey, though at the expense of contamination by an uncertain fraction of asymptotic giant branch (AGB) stars (Robitaille et al. 2008). The surface density of objects from the Robitaille catalogue is shown in Fig. 3 and displays a similar distribution to the RMS YSO sample, with a higher surface density towards the bubbles that sharply drops off to a uniform background level.

The background level is much higher than the RMS YSO sample, as would be expected due to the higher surface density of Robitaille et al. (2008) intrinsically red sources compared to the RMS

<sup>2</sup> <http://www.ast.leeds.ac.uk/RMS>




**Figure 3.** Histogram of the number counts of Robitaille et al. (2008) intrinsically red objects as a function of angular distance from the centre of *Spitzer* bubbles. The distance is expressed in terms of the normalized bubble radius. The number counts are scaled by the area of the annulus corresponding to each bin and thus represent a surface density. Error bars are determined via Poisson statistics.

catalogue (Urquhart et al. 2008). The distribution of Robitaille et al. (2008) objects does not peak at 1 bubble radius, but instead exhibits a relatively flat distribution out to 1 bubble radius. As the RMS catalogue is constrained to star-forming objects at an early evolutionary state (YSOs and UC H II regions), whereas the Robitaille et al. (2008) catalogue is not, this may indicate the presence of an evolutionary gradient across the bubbles. Further classification of the Robitaille et al. (2008) sample and investigation of their star-forming nature are required to prove this hypothesis.

We must also explore the possibility that there may be an intrinsic bias in the distribution of both the RMS and Robitaille et al. (2008) catalogues around the *Spitzer* bubbles due to the common mid-infrared bands used to detect both the bubbles and RMS/Robitaille objects. Although the bubbles are principally identified via their extended PAH emission at 8  $\mu$ m and the RMS YSOs and Robitaille intrinsically red sources are predominantly point infrared sources, the complex mid-infrared environments of the bubbles may lead to a bias in the identification of point sources at their rims. We investigate this possibility by examining the distribution of 6.7 GHz methanol masers drawn from the Methanol MultiBeam (MMB) survey (Green et al. 2009) around the *Spitzer* bubbles. 6.7 GHz methanol masers are thought to exclusively trace young sites of massive star formation (e.g. Menten 1991), and thus allow us to trace the distribution of massive YSOs around the bubbles independent of their mid-infrared emission.

The MMB survey currently occupies a longitude range between  $l = 186$  and  $20$ , i.e. excluding the range  $20 \leq l \leq 186$ , and so only the bubbles in the southern GLIMPSE survey region are presently covered by the MMB survey. The individual masers in the MMB catalogue have had their positions interferometrically determined to sub-arcsecond precision and the maser detections are reported in Caswell (2009), Caswell et al. (2010, 2011), Green et al. (2009, 2010) and Green et al. (in preparation). We plot the surface density



**Figure 4.** Histogram of the number counts of MMB 6.7-GHz masers as a function of angular distance from the centre of *Spitzer* bubbles. The distance is expressed in terms of the normalized bubble radius. The number counts are scaled by the area of the annulus corresponding to each bin and thus represent a surface density. Error bars are determined via Poisson statistics.

of 6.7 GHz MMB masers around the southern *Spitzer* bubbles in Fig. 4.

Fig. 4 displays a distribution of masers very similar to that of RMS YSOs and Robitaille et al. (2008) intrinsically red sources, albeit with larger error bars due to the smaller sample size. There is a clearly distinguished peak in the maser distribution at an angular offset of 1 bubble radius and the surface density of 6.7 GHz masers drops to a roughly constant background level beyond an offset of 2 bubble radii. The peak in the surface density of 6.7 GHz masers appears to be broader than that in the surface density of RMS YSOs; however, the lower signal-to-noise ratio of the MMB surface density makes it difficult to interpret this difference as a real effect.

All three independently selected YSO catalogues (RMS, Robitaille et al. 2008 red sources and MMB 6.7 GHz masers) display very similar surface density distributions and we thus conclude that the increase in surface density of YSOs towards the bubble rims is a real effect. Given this similar behaviour between catalogues, and the currently more comprehensive knowledge of the properties of the RMS YSOs (e.g. Urquhart et al. 2009b,c, 2011; Mottram et al. 2011a), we restrict our further analysis to the RMS YSO catalogue. Although the Robitaille et al. (2008) red source catalogue has greater sample statistics and is likely to be predominantly comprised of YSOs, there is a much greater likelihood of contamination by AGB stars and other non-YSO types than in the RMS YSO catalogue. Similarly, the lower numbers of objects in the MMB 6.7 GHz maser catalogue favours the continuation of our study using the larger and much more studied RMS YSO sample.

## 2.2 The angular cross-correlation of bubbles and YSOs

As a refinement of our simple surface density approach, we also investigated the distribution of RMS YSOs around the bubbles using




an angular two-point cross-correlation analysis, a technique more commonly used to determine the clustering properties of galaxies (e.g. Smith, Boyle & Maddox 1995; Ghirlanda et al. 2006; Bradshaw et al. 2011; Wang et al. 2011). The correlation function defines the probability of finding a population of objects at a particular angular separation from a different second population. Here, we used the RMS YSO sample as our first population ( $D_1$ ) and the Churchwell et al. (2006) catalogue of infrared bubbles for our second population ( $D_2$ ). We calculated the angular cross-correlation using the estimator of Landy & Szalay (1993), modified for the cross-correlation between population 1 and 2 using the equation of Bradshaw et al. (2011), i.e.

$$\omega(\theta) = \frac{N_{D_1 D_2} - N_{D_1 R_2} - N_{R_1 D_2} + N_{R_1 R_2}}{N_{R_1 R_2}}, \quad (1)$$

where  $N_{D_1 D_2}$  represents the normalized number counts at an angular separation of  $\theta$  of RMS source–bubble pairs,  $N_{D_1 R_2}$  and  $N_{R_1 D_2}$  represent the counts of real and random catalogues of RMS source–bubble pairs (and vice versa), and  $N_{R_1 R_2}$  represents the counts between two random catalogues of RMS objects and *Spitzer* bubbles. As in Section 2.1, we scaled  $\theta$  to the radius of the individual *Spitzer* bubble in each pair. To avoid introducing high levels of noise through the randomly generated catalogues, we performed 50 realizations of each catalogue, taking the mean of the results to determine  $\omega(\theta)$ . The errors on  $\omega(\theta)$  were calculated by a bootstrapping approach (e.g. Ghirlanda et al. 2006; Bradshaw et al. 2011). The *Spitzer* bubble catalogue was divided into 100 randomly chosen bootstrap catalogues each matching the original catalogue size (with replacement). The angular cross-correlation was determined for each bootstrap catalogue and the resulting  $1\sigma$  error in  $\omega(\theta)$  is given by the standard deviation in  $\omega(\theta)$  from the 100 random bootstrap samples.

The resulting angular cross-correlation function is plotted in Fig. 5, which reveals almost exactly the same distribution as seen in the surface density distribution shown in Figs 2–4. The RMS YSO sample is found to be strongly correlated with the *Spitzer* bub-



**Figure 5.** The angular cross-correlation of the RMS YSO sample and the catalogue of *Spitzer* bubbles as a function of the normalized bubble radius.

ble catalogue, particularly at the radius corresponding to the rim of the bubbles where the correlation peaks. This peak is significant at the  $9\sigma$  level. The cross-correlation drops sharply beyond the peak at 1 bubble radius and beyond a distance of 2 bubble radii the cross-correlation decreases to essentially zero. This indicates that the probability of finding an RMS YSO near a *Spitzer* bubble is markedly greater at an angular radius of 1 bubble radii, and that beyond an angular distance of 2 bubble radii the RMS YSO population is essentially uncorrelated with the presence of a *Spitzer* bubble.

### 2.3 The angular autocorrelation of RMS YSOs

Finally, we investigate the clustering within the RMS YSO sample, in order to determine whether the previous results in Sections 2.1 and 2.2 are simply due to an intrinsic angular clustering scale within the RMS catalogue that happens to correspond to the typical angular size of a *Spitzer* bubble. The median bubble radius of all the bubbles listed in the Churchwell et al. (2006) catalogue is 1.1 arcmin, with a large observed range stretching between 0.14 and 14 arcmin. However, few bubbles possess extreme radii and two-thirds of the sample have radii between 1 and 3 arcmin. If the RMS YSO sample is naturally clustered on this typical angular scale, then the intrinsic clustering may mimic the apparent overdensity of YSOs observed in the surface density histogram and angular cross-correlation plot (Figs 2 and 5).


To compare the clustering of the control sample to that of the RMS YSO sample associated with bubbles, we calculate the autocorrelation of the sample as a function of angular distance, using the estimator of Landy & Szalay (1993), i.e.

$$\omega(\theta) = \frac{N_{DD} - 2N_{DR} + N_{RR}}{N_{RR}}, \quad (2)$$

where  $N_{DD}$ ,  $N_{DR}$  and  $N_{RR}$  represent the normalized number counts of data–data, data–random and random–random pairs, respectively. We calculate the autocorrelation for three separate samples: the entire RMS YSO sample, those associated with *Spitzer* bubbles (i.e. lying within 2 bubble radii of a particular bubble), and those not associated with any *Spitzer* bubbles (i.e. lying more than 3 bubble radii from *all* bubbles). For simplicity, we respectively refer to these samples as the full RMS YSO sample, the bubble-associated YSO sample and the control sample.

We chose a value of 2 bubble radii for the radius of association due to the steep fall-off in surface density and angular cross-correlation beyond this radius. A total of 116 RMS YSOs are found within 2 bubble radii of a *Spitzer* bubble and 629 RMS YSOs are found at an angular distance greater than 3 bubble radii from any bubble. As in Section 2.2, we constructed random catalogues of each of the three samples and performed 50 realizations of each random catalogue to avoid introducing higher levels of noise.

The autocorrelations for these three samples of RMS YSOs are shown in Fig. 6. The behaviour of the full RMS YSO sample (solid dots in Fig. 6) shows a classic peak towards smaller angular scales and a decrease towards larger angular scales. This implies that the full RMS YSO sample is strongly clustered on scales of  $\sim 1$  arcmin or less. The bubble-associated YSO sample displays a markedly different behaviour, being *anticorrelated* on all angular scales except for a small positive correlation at 2 arcmin. Finally, the control sample shows a correlation on small angular scales similar to the full sample, although much weaker and with a flatter fall-off to large angular scales than the full RMS YSO sample. Interestingly, there is a minor peak in the autocorrelation function at 2 arcmin, though this is not a statistically significant detection. At angular



**Figure 6.** The angular autocorrelation function of RMS YSOs for the full sample of RMS YSOs (solid dots), a ‘control’ sample of RMS YSOs that lie more than 3 bubble radii from all of the Churchwell et al. (2006) bubbles (open squares) and the sample of RMS YSOs that lie within 2 bubble radii of the Churchwell et al. (2006) bubbles (open circles).  $1\sigma$  error bars are shown and are calculated using the bootstrap replacement method.

scales  $\geq 2$  arcmin, the autocorrelation functions of the full RMS YSO sample and the control sample are identical.

Clearly, the autocorrelation function of the bubble-associated RMS YSOs is very different from the other two samples. This implies that the peaks in the YSO surface density and the YSO-bubble angular cross-correlation seen towards the bubble rims are not due to intrinsic clustering within the full RMS YSO sample. The YSOs in the full and control samples are much more highly correlated (i.e. clustered) on smaller angular scales than the bubble-associated YSO sample. On the majority of angles the bubble-associated sample is anticorrelated, which implies that there is a shortfall in the numbers of YSOs associated with the bubbles at these angular separations compared to a random sample. The exception to this is at an angular scale of 2 arcmin, which corresponds closely with the median bubble diameter of 2.2 arcmin. This positive correlation suggests that we may be seeing the signature of RMS YSOs located on either side of the bubble, and the anticorrelation implies that the YSOs are not found on angular scales smaller or larger than this.

We do see a minor secondary peak at 2 arcmin in the autocorrelation functions of the full and control YSO samples, which at first sight suggests that a fraction of these YSOs are correlated at the same angular scale as the median bubble diameter. We expect this behaviour in the full YSO sample (which obviously includes YSOs associated with the bubbles). However, due to the incompleteness of the Churchwell et al. (2006) catalogue caused by the manual search procedure that was used in its construction, we cannot exclude the presence of contaminating bubbles in our control sample. Thus, the autocorrelations that we measure may be artificially enhanced for the control sample, for example at the secondary peak at 2 arcmin. However, this peak is not statistically significant and the full and control samples are clearly more strongly correlated at small scales

than the bubble-associated sample. This suggests that the intrinsic clustering of RMS YSOs does not produce the enhancement of YSOs projected against the rims of the *Spitzer* bubbles. Further work on the much more complete Milky Way Project bubble sample (Simpson et al., in preparation) would aid this analysis.

### 3 DISCUSSION

In the following we combine our results on the YSO surface density distributions, the YSO-bubble angular cross-correlation and the YSO angular autocorrelation together with the measured properties of the bubbles and RMS YSOs. Our particular aims are to investigate the star formation environment of the bubbles in order to determine whether the YSO population is significantly enhanced near the bubbles and if there are any discernible differences in the population of YSOs found near the bubbles when compared to the entire sample. Ultimately, we would like to place statistical constraints on the star formation associated with the bubbles that can inform current and future models of triggered star formation.

#### 3.1 The properties of *Spitzer* bubbles associated with RMS YSOs

Taking the radius of association between YSO and bubbles to be 2 bubble radii, we find that a total of 116 YSOs and UC H II regions from the RMS YSO sample are associated with bubbles from the Churchwell et al. (2006) catalogue. The number of bubbles from the Churchwell et al. (2006) catalogue associated with one or more RMS YSOs is 72, which corresponds to  $22 \pm 3$  per cent of the bubble catalogue (where the error is calculated using Poisson statistics). Of the 72 bubbles associated with RMS YSOs, 30 bubbles are associated with more than one RMS YSO or UC H II, ranging from 2 to 5 objects per bubble (and with a mean of 2.6 objects per bubble for the multiple matches).

The majority of the Churchwell et al. (2006) bubbles are not associated with RMS YSOs. This does not imply that the remaining bubbles are devoid of surrounding star formation, but merely that any star formation which is present is of sufficiently low luminosity to fall below the RMS detection threshold (which is typically  $\geq 1000 L_{\odot}$  at distances of a few kpc; Urquhart et al. 2011). Unfortunately, the majority of the Churchwell et al. (2006) bubbles do not have measured distances that are free from the kinematic distance ambiguity (Deharveng et al. 2010) and so we cannot determine the individual completeness limit for each bubble. However, the typical distances of the bubbles range from 2 to 13 kpc (Deharveng et al. 2010) and so we can confidently say that the present RMS study of bubbles is complete to YSOs with a luminosity of  $\geq 10^4 L_{\odot}$  for the most distant bubbles in the sample (Urquhart et al. 2011) and to YSOs with a luminosity of  $\geq 1000 L_{\odot}$  for bubbles located at the typical distance of a few kpc.

As the luminosity of a B3 star is  $\sim 550 L_{\odot}$  (Meynet & Maeder 2000), we thus identify *massive* star formation associated with the bubbles. Hence,  $22 \pm 3$  per cent of the bubbles are associated with massive star formation in the massive YSO or UC H II region phase. This is a fraction similar to that found by Deharveng et al. (2010) and Watson, Hanspal & Mengistu (2010), who respectively found 18 and 20 per cent of their bubble samples to be associated with either UC H II regions, YSOs or 6.7 GHz methanol masers. The remaining  $78 \pm 3$  per cent of the bubbles may be associated with low to intermediate-mass star formation, but without a more sensitive survey we cannot confirm this hypothesis. The currently underway *Herschel* Hi-GAL survey of the Galactic plane (Molinari et al.

2010a,b) will provide such a sensitive survey of the entire bubble sample and it would be advantageous to revisit the bubbles with the more sensitive Hi-GAL data when it is available.

We searched for differences in the properties of the bubbles that are associated with RMS YSOs and those that are not, in order to try and identify differences in the properties of bubbles that are associated with massive star formation and those that are not. The two most important measured properties of each bubble are the size (strictly the mean radius of the bubbles) and thickness of the diffuse mid-infrared emission comprising the bubble. For both of these properties, we determined the mean values for the 22 per cent of bubbles that are associated with at least one RMS YSO (i.e. within 2 bubble radii) and the remaining 78 per cent of bubbles that are not associated with an RMS YSO.

The mean radius of bubbles that are associated with an RMS YSO is  $3.6 \pm 0.4$  arcmin, compared to the mean radius of unassociated bubbles of  $4.6 \pm 0.3$  arcmin. The mean thickness of bubbles that are associated with RMS YSOs is  $0.92 \pm 0.08$  arcmin, again compared to the unassociated bubble thickness which is  $1.18 \pm 0.07$  arcmin. Quoted errors are the standard error on the mean. In order to determine the significance of these differences in mean radius and thickness, we performed a two-sample unequal variance (heteroscedastic) *t*-test on each pair of samples (RMS YSO-associated bubbles and unassociated bubbles). The *t*-tests return probabilities of 1 and 0.8 per cent, respectively, that the mean radius and thickness of YSO-associated bubbles and unassociated bubbles are drawn from populations with the same mean.

The bubbles that are associated with massive star formation thus tend to be both smaller and thinner than those bubbles that are not associated with massive star formation. These results are not significant at a level of  $3\sigma$  or greater and, combined with the potential biases in the bubble population discussed earlier, should be interpreted with caution. However, it is instructive to speculate on what may be the physical causes behind these observed differences. Weaver et al. (1977) predict that the radius of a wind-blown bubble should increase much faster than the thickness of the swept-up shell when the bubbles are in their first expansion stage. Hence, bubbles with small radii and thinner shells should be younger than larger bubbles with thicker shells, as also suggested by Dale et al. (2009). Of course, we do not have physical distances for the majority of our sample and are thus dealing with angular radii rather than physical radii. Hence, the bubbles that we have identified as small in angular size may just be the more distant members of the sample. However, over the whole sample of bubbles, these effects should average out and our tentative results suggest that it is the younger bubbles within the sample that are more likely to be associated with massive star formation.

Theoretical models of shell fragmentation (the collect and collapse process; Whitworth et al. 1994; Dale, Bonnell & Whitworth 2007) suggest that fragmentation of the swept-up shell of a bubble tends to occur on time-scales of 1 to a few Myr. Very little detailed study of the bubble lifetimes has currently been made. The few bubbles that have been studied to date have dynamical lifetimes of 0.5 to a few Myr (Watson et al. 2008, 2009) and many of the samples are likely to be H II regions powered by late O to early B stars (Bania et al. 2010; Beaumont & Williams 2010; Deharveng et al. 2010; Anderson et al. 2011), which again have main-sequence lifetimes of the order of a few Myr to a few tens of Myr. So the picture that massive star formation tends to be associated with smaller and younger bubbles is largely consistent with the predictions of the collect and collapse models that star formation should happen on  $\sim$  Myr time-scales. However, if these bubbles are shown to be

much younger than 0.5 Myr, then this would point towards the star formation being caused by the implosion of pre-existing density structures (e.g. the EDGI proposed by Walch et al. 2011), or perhaps simply pre-existing untriggered star formation as suggested by Dale & Bonnell (2011).

In the following subsection, we concentrate upon the surface density of YSOs around the bubbles, with the aim of showing that there is a statistically significant overdensity of YSOs and that it is unlikely that the majority of the YSOs formed spontaneously.

### 3.2 An overdensity of YSOs around *Spitzer* bubbles

It is clear from the surface density plots shown in Figs 2–4 that there is a significantly enhanced surface density of YSOs found at the rim of the bubbles. For the RMS YSO sample shown in Fig. 2, the YSO surface density is  $19.1 \pm 3.6$  YSOs per unit area at an angular distance from the bubble centre of 1 bubble radius, some  $4\sigma$  above the mean surface density of YSOs at angular distances greater than 2 bubble radii ( $3.1 \pm 0.2$  YSOs per unit area). This result is confirmed by the angular cross-correlation function of the RMS YSO sample shown in Fig. 5, where the cross-correlation function peaks at an angular distance of 1 bubble radius with a significance of  $9\sigma$ . In the immediate environment of a *Spitzer* bubble, the highest probability location to find an RMS YSO is projected against the rim of the bubble.

Moreover, it is clear from inspecting Fig. 2 that the surface density of YSOs is not only enhanced at an angular offset of 1 bubble radius, but that it is enhanced over the entire angular scale of the bubbles out to an angular offset of 2 bubble radii. We can see this by comparing the mean surface density of YSOs ‘inside’ 2 bubble radii and ‘outside’ 2 bubble radii. The mean surface density of YSOs within an angular offset of 2 bubble radii is  $8.9 \pm 1.7$  YSOs per unit area compared to a value of  $3.2 \pm 0.2$  YSOs per unit area at an angular offset of 2 bubble radii or greater. A two-sample unequal variance (heteroscedastic) *t*-test of these two subsamples returns a probability of only 0.4 per cent that these two subsamples are drawn from populations with the same mean. Hence, we have demonstrated that there is a statistically significant overdensity of massive YSOs associated with the bubbles compared to the background, with an enhanced probability of finding these YSOs projected against the rim of the bubbles.

What do these results imply? First, there is a greater concentration of massive star formation towards the bubbles than in the wider environment. This result is confirmed by the surface density of MMB 6.7 GHz masers (see Fig. 4), which trace a YSO population independent of mid-infrared emission. A greater concentration of star formation towards the bubbles implies that the bubbles are either efficient at producing YSOs, or that they are found in regions of high YSO surface density. This is the classic chicken-and-egg scenario applied to massive star formation: do the bubbles precede the high surface density of YSOs, or does the high surface density of YSOs precede (or occur simultaneously with) the formation of the bubbles?

Before considering this question more fully, we must bring in the second of our results – that there is an enhanced probability of finding YSOs projected against the rim of the bubbles (i.e. at an angular offset of 1 bubble radius). By inspecting the autocorrelation of the RMS YSOs, we showed in Section 2.3 that this effect is not likely to be due to intrinsic clustering within the RMS sample on angular scales similar to the bubble radii. The ancillary question raised by this result is: why are the YSOs more likely to be found

projected against the rim of the bubbles, i.e. what is special about the bubble rims?

The bubble rims are traced by 8  $\mu\text{m}$  PAH emission which originates from the photon-dominated region between the ionization front being driven out by the  $\text{H II}$  region within the bubble and the surrounding neutral medium. The rim of the bubbles thus shows the interface between the  $\text{H II}$  region and surrounding neutral gas. For a spherical bubble morphology, one would expect the column density of gas to be greater at the bubble rims due to the greater path length through the neutral material towards the rims. So, at first glance, the higher surface density of YSOs projected against the bubble rims may simply reflect the higher column density at the rim of the bubbles, i.e. the YSOs trace molecular column density.

However, while the sample of bubbles that have been observed at relatively high angular resolution in molecular lines (Beaumont & Williams 2010) does show a peaked molecular column density profile at a normalized bubble radius of 1, the column density falls off much less sharply than the YSO surface density. An inspection of fig. 2 of Beaumont & Williams (2010) shows that at a normalized bubble radius of 1.5 the CO intensity can be roughly half of that at a normalized radius of 1. This suggests that the YSOs may not trace the column density distribution, although much closer scrutiny of the bubbles in a non-optically thick tracer is required to confirm this hypothesis. Moreover, the CO contrast between the centre of the bubbles and their rims is often extreme (Beaumont & Williams 2010) whereas the YSO surface density within an angular offset of 2 bubble radii is everywhere higher than the background level. Thus, we cannot confidently say that the YSO surface density traces the gas column density around the bubbles.

The YSO surface density is strongly peaked at an offset of 1 bubble radius and decreases sharply beyond this value. Beyond an angular offset of 2 bubble radii, the surface density of YSOs is essentially undistinguishable from the background level. The angular cross-correlation function shows a similar steep drop – beyond an angular distance of 2 bubble radii, the bubbles and RMS YSOs are essentially *uncorrelated*. The implication of this is that whatever causes the rise in YSO surface density is closely related to the rim of the bubbles. The bubble radius is a dynamic value and expected to increase over time as stellar winds or radiation pressure causes the bubbles to expand. Combined with this is the fact that the massive YSOs and UC  $\text{H II}$  regions identified by the RMS survey typically tend to have lifetimes around a few  $10^4$ – $10^5$  years (Mottram et al. 2011b) and so should trace very recent star formation.

The sum of these pieces of information leads us to conclude that it is likely that the bubbles predate the YSOs. If the bubbles formed in an environment with a high surface density of YSOs (e.g. in the turbulent highly fragmented initial conditions suggested by Dale & Bonnell 2011), then the distribution should not peak at the rim of the bubble as the bubble radius is time dependent. Similar arguments have been used by Preibisch et al. (2011) for YSOs detected at the edges of shells in Carina. Also, in this case the extent of the enhanced YSO surface density should also not be related to the current radius of the bubble – why are bubbles found in regions of enhanced YSO surface density occupying twice their *current* angular radius? Finally, the relative time-scales of the massive YSOs and those required for the expansion of the bubbles imply that the YSOs formed *after* the bubbles. We thus conclude that a significant fraction of the YSOs seen against the rim of the bubbles were likely triggered by the expansion of the bubble.


A greater understanding of the dynamical time-scales for the expansion of the bubbles and also the molecular environment of

the bubbles are required to confirm this hypothesis. Pinpointing the YSO formation to have occurred after the bubble was formed is crucial to disentangling cause and effect in the star formation surrounding the bubbles. Currently, only a few bubbles have had their dynamical lifetimes estimated and more studies similar to those of Watson et al. (2009) are required over a larger sample of bubbles. Comparing the YSO distribution to the gas distribution is also crucial to investigate differences in the population of YSOs at the rims of *Spitzer* bubbles, for example to determine whether the star formation efficiency is enhanced at the bubble rims. Finally, it will also be instructive to apply the same statistical tools that we have used in this paper to the latest generation of triggered star formation models (e.g. Dale & Bonnell 2011; Walch et al. 2011) in order to see whether the surface density of sink particles in the SPH simulations matches that of YSOs around observed bubbles.

### 3.3 The luminosity function of RMS YSOs associated with bubbles


The RMS survey has determined luminosities for their entire sample of YSOs and UC  $\text{H II}$  regions (Mottram et al. 2011a; Urquhart et al. 2011), and so in this subsection we seek to identify differences in the luminosity function between those YSOs that are associated with bubbles (i.e. within an angular offset of 2 bubble radii) compared to the full population. As both luminosity functions are essentially power laws with turnovers caused by incompleteness at low luminosities, one must take care that any differences between the two samples are not primarily due to differences in the completeness of each sample. To avoid this issue, we cut both samples at a luminosity of  $10^4 L_\odot$ , i.e. the minimum YSO luminosity at which the RMS catalogue is complete (Urquhart et al. 2011). We plot these truncated luminosity functions in Fig. 7 and the corresponding cumulative distribution functions in Fig. 8.

Inspecting Fig. 7 shows that the RMS YSOs associated with the bubbles do not tend to have a higher luminosity than the rest of the



**Figure 7.** The luminosity function of the entire RMS YSO sample (solid line) and those YSOs that lie within an angular offset of 2 bubble radii from a *Spitzer* bubble (dashed line).





**Figure 8.** The cumulative distribution function of the luminosities of the entire RMS YSO sample (solid line) and those YSOs that lie within an angular offset of 2 bubble radii from a *Spitzer* bubble (dashed line).

population. The two luminosity functions are essentially identical. A Kolmogorov–Smirnov test on the two luminosity functions yields the result that there is a 27 per cent probability that the two distributions are drawn from the same sample, and thus we cannot reject the null hypothesis with any significance. If the YSOs associated with the bubbles were triggered (as we argue in Section 3.2), then we conclude that the triggering process does not result in stars with an appreciably different luminosity function.

This result must be contrasted against the many inferences that have been made to date which suggest that the triggering process may produce stars with a luminosity function skewed to higher luminosities (e.g. Sugitani et al. 1989, 1991; Dobashi et al. 2001; Deharveng, Zavagno & Caplan 2005; Motoyama, Umemoto & Shang 2007; Urquhart, Morgan & Thompson 2009a). Fragmentation models for swept-up shells around massive stars predict a top-heavy mass distribution of fragments, which in turn may lead to a top-heavy initial mass function (IMF) distribution (Whitworth et al. 1994; Dale et al. 2009). Observations of molecular clouds associated with H II regions also suggested that *IRAS* point sources associated with the clouds nearer the H II regions are more luminous than those found to be more distant (Yamaguchi et al. 1999). However, these results are not consistent with the largely constant form of the IMF observed over Galactic scales. If the triggering process does induce a top-heavy IMF, then the results of triggering do not dominate the IMF (Dale et al. 2009). Here we have shown that if these YSOs are triggered, then the triggering process results in a luminosity function that is indistinguishable from the full YSO sample.

### 3.4 The fraction of massive stars in the Milky Way that may have been triggered

Estimating the impact of the triggering process in global Galactic star formation is a vital part of determining a simple prescription for star formation that can be applied to galaxy evolution models.

Because we have combined the results of two large area and relatively unbiased surveys (the RMS survey and the Churchwell et al. 2006 bubble catalogue), we are in a position to try and estimate the contribution of triggered star formation to the Galaxy’s population of stars. Before doing so, we must stress the major caveat involved – that the Churchwell et al. (2006) bubble catalogue is likely to be highly incomplete. Thus, we do not attempt a detailed treatment at this stage and simply infer a lower limit to the fraction of massive stars in the Milky Way that *may* have been triggered.

As discussed in Section 3.1 we find 116 YSOs and UC H II regions from the RMS YSO sample that are associated with bubbles from the Churchwell et al. (2006) catalogue. The bubble-associated YSOs form 14 per cent of the 846 objects contained in the RMS YSO sample over the GLIMPSE survey region. The RMS survey is complete for essentially all massive YSOs with luminosities in excess of  $10^4 L_{\odot}$  out to the furthest kinematic distance in the bubble sample ( $\sim 14$  kpc). Thus, assuming that the incompleteness of the Churchwell catalogue dominates over the fraction of the bubble-associated YSOs that were triggered, we estimate that at least 14 per cent of the massive stars in the Milky Way could have been triggered. If, as suggested by Churchwell et al. (2006), the *Spitzer* bubble catalogue is  $\sim 50$  per cent incomplete then the true fraction of triggered massive stars could be up to  $\sim 30$  per cent.

## 4 SUMMARY AND CONCLUSIONS

We have carried out a detailed statistical study of YSOs found nearby *Spitzer* bubbles from the Churchwell et al. (2006) catalogue. Our main results are summarized below.

(i) The surface densities of YSOs and UC H II regions from the RMS survey (Urquhart et al. 2011), intrinsically red sources from Robitaille et al. (2008) and MMB 6.7 GHz methanol masers (Green et al. 2009) are enhanced towards *Spitzer* bubbles from the Churchwell et al. (2006) catalogue. The surface density of all three YSO catalogues peaks towards the projected angular radius of the bubbles, with a peak surface density of  $4\sigma$  above the mean background level for RMS YSOs and UC H II regions. The mean surface density of YSOs associated with the bubbles is overdense with respect to the surrounding mean background at the  $3\sigma$  level.

(ii) The angular cross-correlation function of RMS YSOs and *Spitzer* bubbles shows a behaviour similar to the surface density, with a  $9\sigma$  peak in the cross-correlation function at an angular offset of 1 bubble radius. In the immediate environment of a *Spitzer* bubble, the highest probability location to find an RMS YSO is projected against the rim of the bubble. RMS YSOs and *Spitzer* bubbles are essentially uncorrelated beyond an angular distance of 2 bubble radii. Examination of the autocorrelation functions suggests that these effects are not caused by intrinsic clustering within the RMS YSO sample.

(iii) Most *Spitzer* bubbles are not associated with massive YSOs:  $22 \pm 3$  per cent of the Churchwell catalogue are associated with RMS YSOs. This fraction is consistent with smaller studies (Deharveng et al. 2010; Watson et al. 2010).

(iv) *Spitzer* bubbles associated with RMS YSOs tend to possess both thinner rims and smaller angular radii than bubbles that are not associated with RMS YSOs. We interpret this tendency to be due to an age effect, with RMS YSOs forming around younger and smaller bubbles.

(v) The different relative time-scales for the formation of *Spitzer* bubbles and RMS YSOs, and the strong peak in surface density and cross-correlation at the rims of the bubbles, lead us to conclude

that a significant fraction of the RMS YSOs were triggered by the expansion of the bubble.

(vi) We find no significant differences in the luminosity function of RMS YSOs associated with *Spitzer* bubbles compared to the entire RMS YSO population, which suggests that the triggering process does not result in a top-heavy luminosity function or IMF.

(vii) We estimate from the fraction of RMS YSOs associated with bubbles and the incompleteness of the bubble catalogue that the lower limit for the fraction of massive stars in the Milky Way that could have been triggered is 14 per cent. If the Churchwell et al. (2006) catalogue is 50 per cent incomplete then the upper limit to the fraction of massive stars that could have been triggered may be up to  $\sim 30$  per cent. Therefore, this mode of massive star formation ought not to be ignored when considering star formation on Galactic scales.

We must stress that these results are based on the current and largely incomplete sample of known *Spitzer* bubbles, which were identified by manual searches of the GLIMPSE image data base (Churchwell et al. 2006). However, the groundwork for a much more comprehensive catalogue of bubbles is currently being laid by the Milky Way Project, which aims to identify many more bubbles by a systematic citizen science survey of the GLIMPSE images (Simpson et al., in preparation). In addition, the *Herschel* Hi-GAL survey (Molinari et al. 2010b) holds the promise of a much more complete YSO catalogue reaching fainter luminosities than the RMS survey.

Statistical studies of the type that we have presented here will be of increasing importance in the age of large-scale surveys of the Milky Way, and offer the prospect of being able to directly compare the observed distribution of YSOs with the predicted distribution of sink particles in triggered star formation models (e.g. Dale & Bonnell 2011; Walch et al. 2011). With these advances, the study of triggered star formation will finally move beyond the phenomenological stage to be able to make direct predictions.

## ACKNOWLEDGMENTS

We would like to thank the anonymous referee for a report that was both prompt and illuminating. We thank Steve Longmore and Willem Baan for useful and illuminating discussions on potential biases in YSO identification and the incompleteness of the bubble catalogue. We would also like to thank Andrew Walsh for organizing a stimulating meeting of observers and theorists in Townsville that provided the initial seed for this work. This research would not have been possible without the NASA Astrophysics Data System Bibliographic Services. This paper made use of information from the Red MSX Source survey data base at [www.ast.leeds.ac.uk/RMS](http://www.ast.leeds.ac.uk/RMS), which was constructed with support from the Science and Technology Facilities Council of the UK. We made use of positional data on 6.7 GHz methanol masers provided by the Methanol Multi-Beam (MMB) survey, also supported by the Science and Technology Facilities Council, and with a maser data base available at <http://astromasers.org>.

## REFERENCES

Anderson L. D., Bania T. M., Balser D. S., Rood R. T., 2011, *ApJS*, 194, 32  
 Bania T. M., Anderson L. D., Balser D. S., Rood R. T., 2010, *ApJ*, 718, L106  
 Bastian N., Gieles M., Efremov Y. N., Lamers H. J. G. L. M., 2005, *A&A*, 443, 79  
 Beaumont C. N., Williams J. P., 2010, *ApJ*, 709, 791  
 Benjamin R. A. et al., 2003, *PASP*, 115, 953

Bertoldi F., 1989, *ApJ*, 346, 735  
 Bisbas T. G., Wunsch R., Whitworth A. P., Hubber D. A., 2009, *A&A*, 497, 649  
 Bisbas T. G., Wunsch R., Whitworth A. P., Hubber D. A., Walch S., 2011, arXiv e-prints  
 Bradshaw E. J. et al., 2011, *MNRAS*, 415, 2626  
 Carey S. J. et al., 2009, *PASP*, 121, 76  
 Caswell J. L., 2009, *Publ. Astron. Soc. Aust.*, 26, 454  
 Caswell J. L. et al., 2010, *MNRAS*, 404, 1029  
 Caswell J. L. et al., 2011, *MNRAS*, 417, 1964  
 Churchwell E. et al., 2006, *ApJ*, 649, 759  
 Churchwell E. et al., 2007, *ApJ*, 670, 428  
 Dale J. E., Bonnell I., 2011, *MNRAS*, 414, 321  
 Dale J. E., Bonnell I. A., Whitworth A. P., 2007, *MNRAS*, 375, 1291  
 Dale J. E., Wunsch R., Whitworth A., Palouš J., 2009, *MNRAS*, 398, 1537  
 Deharveng L., Zavagno A., Caplan J., 2005, *A&A*, 433, 565  
 Deharveng L. et al., 2010, *A&A*, 523, A6  
 Dobashi K., Yonekura Y., Matsumoto T., Momose M., Sato F., Bernard J.-P., Ogawa H., 2001, *PASJ*, 53, 85  
 Elmegreen B. G., 2011, *EAS Publ. Ser.*, 51, 45  
 Elmegreen B. G., Lada C. J., 1977, *ApJ*, 214, 725  
 Ghirlanda G., Magliocchetti M., Ghisellini G., Guzzo L., 2006, *MNRAS*, 368, L20  
 Green J. A. et al., 2009, *MNRAS*, 392, 783  
 Green J. A. et al., 2010, *MNRAS*, 409, 913  
 Landy S. D., Szalay A. S., 1993, *ApJ*, 412, 64  
 Lefloch B., Lazareff B., 1994, *A&A*, 289, 559  
 Menten K. M., 1991, *ApJ*, 380, L75  
 Meynet G., Maeder A., 2000, *A&A*, 361, 101  
 Miao J., White G. J., Thompson M. A., Nelson R. P., 2009, *ApJ*, 692, 382  
 Molinari S. et al., 2010a, *A&A*, 518, L100  
 Molinari S. et al., 2010b, *PASP*, 122, 314  
 Morgan L. K., Thompson M. A., Urquhart J. S., White G. J., Miao J., 2004, *A&A*, 426, 535  
 Morgan L. K., Thompson M. A., Urquhart J. S., White G. J., 2008, *A&A*, 477, 557  
 Motoyama K., Umemoto T., Shang H., 2007, *A&A*, 467, 657  
 Mottram J. C. et al., 2011a, *ApJ*, 730, L33  
 Mottram J. C. et al., 2011b, *A&A*, 525, A149  
 Oey M. S., Watson A. M., Kern K., Walth G. L., 2005, *AJ*, 129, 393  
 Pomarès M. et al., 2009, *A&A*, 494, 987  
 Preibisch T. et al., 2011, *A&A*, 530, A34  
 Robitaille T. P. et al., 2008, *AJ*, 136, 2413  
 Smith R. J., Boyle B. J., Maddox S. J., 1995, *MNRAS*, 277, 270  
 Sugitani K., Ogura K., 1994, *ApJS*, 92, 163  
 Sugitani K., Fukui Y., Mizuni A., Ohashi N., 1989, *ApJ*, 342, L87  
 Sugitani K., Fukui Y., Ogura K., 1991, *ApJS*, 77, 59  
 Sugitani K., Tamura M., Ogura K., 1999, in Nakamoto T., ed., *Star Formation 1999. Nobeyama Radio Observatory*, p. 358  
 Sugitani K., Matsuo H., Nakano M., Tamura M., Ogura K., 2000, *AJ*, 119, 323  
 Thompson M. A., White G. J., Morgan L. K., Miao J., Fridlund C. V. M., Hultgren-White M., 2004, *A&A*, 414, 1017  
 Urquhart J. S., Thompson M. A., Morgan L. K., White G. J., 2004, *A&A*, 428, 723  
 Urquhart J. S., Thompson M. A., Morgan L. K., White G. J., 2006, *A&A*, 450, 625  
 Urquhart J. S., Hoare M. G., Lumsden S. L., Oudmaijer R. D., Moore T. J. T., 2008, in Beuther H., Linz H., Henning T., eds, *ASP Conf. Ser. Vol. 387, Massive Star Formation: Observations Confront Theory*. Astron. Soc. Pac., San Francisco, p. 381  
 Urquhart J. S., Morgan L. K., Thompson M. A., 2009a, *A&A*, 497, 789  
 Urquhart J. S. et al., 2009b, *A&A*, 501, 539  
 Urquhart J. S. et al., 2009c, *A&A*, 507, 795  
 Urquhart J. S. et al., 2011, *MNRAS*, 410, 1237  
 Walch S., Whitworth A., Bisbas T., Hubber D. A., Wunsch R., 2011, (arXiv:1109.3478)  
 Wang L. et al., 2011, *MNRAS*, 414, 596

- Watson C. et al., 2008, *ApJ*, 681, 1341  
 Watson C., Corn T., Churchwell E. B., Babler B. L., Povich M. S., Meade M. R., Whitney B. A., 2009, *ApJ*, 694, 546  
 Watson C., Hanspal U., Mengistu A., 2010, *ApJ*, 716, 1478  
 Weaver R., McCray R., Castor J., Shapiro P., Moore R., 1977, *ApJ*, 218, 377  
 Whitworth A. P., Francis N., 2002, *MNRAS*, 329, 641  
 Whitworth A. P., Bhattal A. S., Chapman S. J., Disney M. J., Turner J. A., 1994, *A&A*, 290, 421  
 Yamaguchi R., Saito H., Mizuno N., Mine Y., Mizuno A., Ogawa H., Fukui Y., 1999, *PASJ*, 51, 791  
 Zavagno A., Deharveng L., Comerón F., Brand J., Massi F., Caplan J., Russeil D., 2006, *A&A*, 446, 171  
 Zavagno A., Pomarès M., Deharveng L., Hosokawa T., Russeil D., Caplan J., 2007, *A&A*, 472, 835  
 Zavagno A. et al., 2010a, *A&A*, 518, L81  
 Zavagno A. et al., 2010b, *A&A*, 518, L101

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.