

Impact of co-morbid burden on mortality in patients with coronary heart disease, heart failure, and cerebrovascular accident: a systematic review and meta-analysis

Muhammad Rashid^{1*}, Chun Shing Kwok¹, Chris P. Gale², Patrick Doherty³, Ivan Olier¹, Matthew Sperrin⁴, Evangelos Kontopantelis⁴, George Peat⁵, and Mamas A. Mamas^{1,6}

¹Keele Cardiovascular Research Group, Institute for Science and Technology in Medicine, Guy Hilton Research Centre, Keele University, Thornburrow Drive, Hartshill, Stoke-on-Trent ST4 7QB, UK; ²Leeds Institute of Cardiovascular and Metabolic Medicine, University of Leeds, Leeds, UK; ³University of York, York, UK; ⁴Far Institute, Institute of Population Health, University of Manchester, Manchester, UK; ⁵Institute for Primary Care and Health Sciences, University of Keele, Keele, UK; and ⁶Royal Stoke Hospital, University Hospital North Midlands, Stoke-on-Trent, UK

Received 13 March 2016; revised 29 April 2016; accepted 5 May 2016; online publish-ahead-of-print 10 May 2016

Aims

We sought to investigate the prognostic impact of co-morbid burden as defined by the Charlson Co-morbidity Index (CCI) in patients with a range of prevalent cardiovascular diseases.

Methods and results

We searched MEDLINE and EMBASE to identify studies that evaluated the impact of CCI on mortality in patients with cardiovascular disease. A random-effects meta-analysis was undertaken to evaluate the impact of CCI on mortality in patients with coronary heart disease (CHD), heart failure (HF), and cerebrovascular accident (CVA). A total of 11 studies of acute coronary syndrome (ACS), 2 stable coronary disease, 5 percutaneous coronary intervention (PCI), 13 HF, and 4 CVA met the inclusion criteria. An increase in CCI score per point was significantly associated with a greater risk of mortality in patients with ACS [pooled relative risk ratio (RR) 1.33; 95% CI 1.15–1.54], PCI (RR 1.21; 95% CI 1.12–1.31), stable coronary artery disease (RR 1.38; 95% CI 1.29–1.48), and HF (RR 1.21; 95% CI 1.13–1.29), but not CVA. A CCI score of >2 significantly increased the risk of mortality in ACS (RR 2.52; 95% CI 1.58–4.04), PCI (RR 3.36; 95% CI 2.14–5.29), HF (RR 1.76; 95% CI 1.65–1.87), and CVA (RR 3.80; 95% CI 1.20–12.01).

Conclusion

Increasing co-morbid burden as defined by CCI is associated with a significant increase in risk of mortality in patients with underlying CHD, HF, and CVA. CCI provides a simple way of predicting adverse outcomes in patients with cardiovascular disease and should be incorporated into decision-making processes when counselling patients.

Keywords

Charlson Co-morbidity Index • Cardiovascular disease • Mortality

Introduction

Cardiovascular disease (CVD) is a major cause of morbidity and mortality, accounting for 30% of all-cause mortality worldwide.¹ Given the incidence of CVD and co-morbidity burden increases with age,² a significant proportion of patients with CVD are older with multiple co-morbidities. This affects disease progression and clinical outcomes and can influence clinical decision-making.^{3–5}

Cardiovascular co-morbidities such as hypertension, diabetes, atrial fibrillation, heart failure, and stroke have an independent association with increased mortality in patients hospitalized with acute myocardial infarction with increasing numbers of these co-morbidities particularly associated with poor outcomes.⁶

While previous studies have mainly focused on cardiovascular co-morbid conditions, patients with CVD often have a broad spectrum of non-cardiovascular co-morbidities. It remains unclear, however,

* Corresponding author. Tel: +44 01782 671652, Fax: +44 01782 674467, Email: doctorrashid7@gmail.com

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2016. For permissions please email: journals.permissions@oup.com.

Table 1 Charlson co-morbidity index

Variable	Points
Myocardial infarction	1
Congestive heart failure	1
Peripheral vascular disease	1
Cerebrovascular disease	1
Dementia	1
Chronic obstructive pulmonary disease	1
Connective tissue disease	1
Peptic ulcer disease	1
Diabetes mellitus	1 if uncomplicated 2 if end-organ damage
Moderate to severe chronic kidney disease	2
Hemiplegia	2
Leukaemia	2
Malignant lymphoma	2
Solid tumour	2 6 if metastatic
Liver disease	1 if mild 3 if moderate to severe
AIDS	6

how clustering of multiple cardiovascular and or non-cardiovascular chronic conditions influences clinical outcomes. Therefore, there is a need to understand the impact of co-morbid burden, rather than focusing on individual co-morbid conditions on clinical outcomes in patients with prevalent CVD.²

The Charlson Co-morbidity Index (CCI) is a recognized measure of co-morbid burden⁷ and quantifies the prognostic impact of 22 co-morbid conditions based on their number and individual prognostic impact by means of a score.⁸ It is a useful tool for estimating prognosis in patients with multiple coexisting illnesses. Table 1 represents the variables. Although various studies have evaluated the prognostic value of CCI in predicting outcomes in different cohorts of patients with CVD, there is no systematic review of the literature that evaluates the prognostic value of CCI on mortality across a range of CVDs. In this systematic review, we sought to investigate the prevalence, and prognostic impact, of co-morbidity defined by the CCI score in patients with three major CVDs; coronary heart disease (CHD), heart failure, and cerebrovascular accident (CVA).

Methods

Study inclusion criteria

We included primary studies that evaluated the prognostic impact of co-morbid burden defined by CCI in patients with CHD, acute or chronic heart failure, and CVA. Studies were considered for inclusion and detailed review if their abstract potentially met all three of the following criteria:

- (1) Primary studies evaluating the impact of co-morbidity defined by CCI on adverse outcomes in patients with CVD.

- (2) CVD was defined by CHD [comprising of patients undergoing percutaneous coronary intervention (PCI) or stable angina or acute coronary syndrome (ACS)], or acute or chronic heart failure, or cerebrovascular disease.
- (3) Adverse outcomes included mortality and major adverse cardiac events (MACE) at any length of follow-up.

We excluded studies that did not have results on outcomes defined by CCI score, but there was no restriction on the basis of language of study. We also excluded expert opinion and editorial reviews. We included conference abstracts to minimize publication bias.

Search strategy

We searched MEDLINE and EMBASE on July 2015 using the broad search terms: ('Charlson co-morbidity index' OR 'Charlson index' OR 'Charlson co-morbidity score' OR 'Charlson score') AND ('acute myocardial infarction' OR 'acute coronary syndrome' OR 'coronary heart disease' OR 'coronary artery disease' OR 'stroke' OR 'cerebrovascular disease' OR 'cerebrovascular accident' OR 'heart failure' OR 'cardiac failure') AND ('mortality' OR 'death' OR 'major adverse cardiovascular event' OR 'major adverse cardiac event' OR 'cardiovascular disease'). The search results were reviewed by two independent investigators (M.R. and C.S.K.) for studies that met the inclusion criteria, and relevant reviews were identified. Additional studies were retrieved by checking the bibliographies of included studies and relevant reviews.

Data extraction

Data were extracted from each study into preformatted tables generated in Microsoft Word. Data collected included year, country, number of participants, mean age of participants, percentage of male participants, participant inclusion criteria, follow-up assessment, lost to follow-up, and results of association between CCI and outcomes. With regards to quality assessment, we documented the design of the study, reliable method of ascertainment of outcomes, >10% loss to follow-up, and if there was any adjustment for potential confounders.

Data analysis

Meta-analysis for estimated pooled risk ratios (RR) was performed by the inverse variance method using a random-effects model on the software RevMan 5.3 (Nordic Cochrane Centre, København, Denmark). To reduce the risk of confounding associated with crude estimates, where available, we chose to pool the results from the most adjusted model, whereby results were expressed as pooled relative RR with accompanying 95% confidence intervals (CI). Statistical heterogeneity was assessed using the I^2 statistic, with values of 30–60% representing a moderate level of heterogeneity.⁹ For $I^2 > 50\%$, we performed sensitivity analysis by systematic exclusion of studies and evaluated the effect on I^2 estimates (see Supplementary material online, Table S1). The primary analysis evaluated adverse outcomes with incremental increase in CCI, and secondary analysis was performed by considering higher group of CCI score vs. lower group of CCI score. In the final analysis, we excluded studies by the same research group over the same time period where there was the potential that the same participants were studied more than once. Where there were similar study participants, we chose the study with the largest sample size or highest adverse outcome event rate. We evaluated publication bias through Funnel plots and Egger's test where there were >10 studies in the analysis and no evidence of statistical heterogeneity as the power to detect publication bias was low for meta-analyses of 10 or fewer studies.¹⁰

Table 2 Study design and characteristics of participants

Study ID	Study design; Year; Country	No. of Participants	Participants with CCI = 0 (%)	Mean age	% Male	Description of participants
Bottle 2013 ¹¹	Retrospective cohort study; 2006–2009; UK	288 550	15 177 (5%)	42% of admissions > 75 years of age	61%	Participants were emergency admissions for ACS in England
Bar 2011 ¹²	Retrospective cohort study; 2001–2004; USA	243	88 (36%)	NA	NA	Patients with non-traumatic intra-cerebral haemorrhage presented to hospital emergency department
Chin 1998 ¹³	Prospective cohort study; 1993–1994; USA	257	48 (18%)	Full cohort 41% > 70 years of age	Full cohort 47%	Participants were admitted with congestive heart failure to the Brigham and Women's Hospital
Chirinos 2006 ¹⁴	Prospective cohort study; 1998–2000; USA	305	70 (22%)	64 years	100%	Male veterans undergoing coronary angiography at Miami Veterans Administration Medical Centre
Clarke 2011 ¹⁵	Retrospective cohort study; 1998–2004; Canada	824	NA	64 years	69%	Consecutive patients followed at a tertiary care specialty ambulatory heart failure clinic
Eberli 2013 ¹⁶	Prospective registries; NA; international	5559	2041 (36%)	NA	NA	Participants from e-Biomatrix PMR and PMS registries evaluating the efficacy and safety of biolimus-A9-eluting stent
Erickson 2014 ¹⁷	Retrospective cohort study; 1999–2007; USA	1202	NA	64 years	65%	Participants from ACS registry from a large university hospital
Fabbian 2013 ¹⁸	Retrospective cohort study; 1999–2009; Italy	88 014	NA	71 years	48%	Participants from database of Emilia-Romagna region, Italy who presented with first event of myocardial infarction
Goldstein 2004 ¹⁹	Prospective cohort study; 1995–1997; USA	960	212 (22%)	68 years	NA	Participants admitted with ischaemic stroke, Department of Veterans Affairs (VA) Stroke Study
Hong 2011 ²⁰	Prospective cohort study; 2006–2008; international	675	NA	83 years	58%	Octogenarian participants from Sirolimus-eluting coronary stent (e-Select) registry
Huang 2015 ²¹	Retrospective; 2002–2011; Taiwan	798 328	315 556 (39%)	45% ≥ 65 years	57%	Participants with disabilities from the National Health Insurance Research Database published by the Ministry of Health and Welfare in Taiwan
Jeger 2014 ²²	Retrospective; 2005–2012; Switzerland	1909	NA	65 years	78%	Participants from AMIS plus registry
Jong 2002 ²³	Retrospective; 1994–1997; Canada	38 702	15 020 (38%)	85% ≥ 65 years	49%	Participants from Canadian institute for health information database admitted with first diagnosis of heart failure
Khawaja 2014 ²⁴	NA; 2008–2013; USA	383	37 (9%)	NA	NA	Patients with primary intra cerebral haemorrhage
Mamas 2015 ²⁵	Post hoc analysis of prospective registry; 2008–2013; international	3 067	787 (25%)	64 years	78%	Participants were in the Nobori 2 study who underwent Nobori biolimus-eluting stent implantation
Menendez-Colino 2013 ²⁶	NA; Spain	652	NA	85 years	NA	Patients admitted with heart failure in six Spanish hospitals
Munoz-Rivas 2009 ²⁷	Retrospective cohort study; 2005–2007; Spain	270	NA	78 years	42%	Patients with chronic heart failure diagnosis
Núñez 2004 ²⁸	Prospective cohort study; 2000–2003, Spain	1035	481 (46%)	70 years	70%	Patients admitted with diagnosis of acute myocardial infarction
Oudejans 2012 ²⁹	Prospective cohort study; 2003–2007; Netherlands	93	0	83 years	37%	Patients with diagnosis of heart failure

Perez-Barquero 2010 ³⁰	Retrospective cohort study; 2000–2001; Spain	2127	NA	77 years	43%	Patients admitted with heart failure to various hospitals in Spain
Radovanovic 2014 ³¹	Prospective cohort study; 2002–2012; Switzerland	29 620	15 754 (51%)	64 years	73%	Participants from AMIS plus registry
Ramirez-Marrero 2011 ³²	Retrospective cohort study; 2004–2005; Spain	715	NA	66 years	NA	Patients admitted with diagnosis of non-ST-elevation acute coronary syndromes
Ramirez-Marrero 2013 ³³	Retrospective cohort study; 2008–2009; Spain	146	NA	78 years	63%	Patients undergoing percutaneous coronary revascularization
Rodriguez-Pascual 2012 ³⁴	Prospective cohort study; 2006–2009	581	121 (20%)	86 years	33%	Patients admitted to an acute geriatric unit with decompensated heart failure
Sachdev 2004 ³⁵	Prospective cohort study; 1985–1989; USA	1471	810 (55%)	60 years	72%	All patients undergoing initial coronary angiography for symptoms of chronic CAD and found to have significant disease ($\geq 75\%$ stenosis) in one or more coronary arteries
Sanchis 2011 ³⁶	Prospective cohort study; 2002–2009; Spain	1017	NA	68 years	66%	Patients admitted with diagnosis of non-ST-elevation acute coronary syndromes
Schmidt 2012 ³⁷	Retrospective cohort study; 1984–2009; Denmark	234 331	164 937 (70%)	75 years	62%	Patients from nationwide Danish cohort registry admitted with myocardial infarction
Singh 2011 ³⁸	Prospective cohort study; 2005–2008; USA	629	NA	75 years	69%	Patients undergoing PCI at the Mayo Clinic in Rochester, NY, USA
Subramanian 2007 ³⁹	Prospective cohort study; unclear; USA	494	NA	68 years	NA	Participants from Veterans Affairs outpatients with diagnoses of congestive heart failure
Teng 2014 ⁴⁰	Prospective cohort study; 2000–2009; Australia	17 379	105 (0.6%)	70 years	58%	Participants were Aboriginal and non-Aboriginal patient with first heart failure hospitalization
Testa 2009 ⁴¹	Prospective cohort study; 1992–2003; Italy	1268	NA	74 years	43%	Participants from 'Osservatorio Geriatrico Regione Campania' with and without heart failure
Theuns 2011 ⁴²	Prospective; 1999–2008; international	463	NA	62 years	75%	Participants from two ICD registries from Rotterdam and Basel
Tuttolomondo 2008 ⁴³	Retrospective; 1988–1998; Italy	1878	0	77 years	49%	Participants from GIFA registry
Urban 2011 ⁴⁴	Prospective cohort study; 2006–2008; international	15 147	NA	62 years	75%	Participants from Sirolimus-Eluting Coronary Stent implantation study (e-Select) registry
Van Wijk 2013 ⁴⁵	Post hoc analysis of RCT; unclear; international	499	NA	NA	NA	Participants from heart failure study randomized to intensified NT-proBNP-guided vs. symptom-guided therapy

NA, not available or not reported.

Results

Description of included studies

A total of 35^{11–45} studies met the inclusion criteria. The process of study selection is shown in *Figure 1*. The details of the studies’ design and participants are described in *Table 2*. The included studies comprised 14 retrospective cohort studies,^{11,12,15,17,18,21–23,27,30,32,33,37,43} 17 prospective cohort studies,^{13,14,16,19,20,28,29,31,34–36,38–42} 1 *post hoc* analysis of registry,²⁵ and 1 *post hoc* analyses of RCT,⁴⁵ while 2 abstract studies^{24,26} were not clear in reporting the design. There were a total of 1 538 793 participants in 35 studies. Twenty-four studies reported a mean age of 71 years and 62% male. The study size varied from 93 participants³¹ to 798 328 participants.²¹ The follow-up time ranged from 30 days to 5 years.

Seventeen studies^{12–16,19,20,22,23,25,28,29,31,34–36,38,42} reported individual CCI scores and 530 457 out of 1 538 793 (35%) patients had no co-morbidities (CCI = 0). The prevalence of each co-morbid condition in each of the cardiovascular conditions/events studied is presented in *Figure 2*. Diabetes and a history of previous myocardial infarction were the two most common conditions present in patients with CHD. Approximately 10% of the patients with heart failure had previous history of myocardial infarction (only reported in 6 studies out of the total 13) and 12% had a history of chronic obstructive pulmonary disease. Similarly, diabetes was the most prevalent co-morbidity in the patients with CVA cohort. Haematological malignancies such as lymphoma leukaemia and AIDS were the least frequent co-morbid conditions across all the cohorts studied.

Quality assessment of included studies


The quality of studies included is described in *Table 3*. There was no loss to follow-up for 13 of the included studies. Twenty-two studies had <10% loss to follow-up. The largest absolute loss to follow-up was reported by Radovanovic *et al.* as they excluded 1091 patients from final results due to unavailability of CCI data.³¹ Just over half of the studies^{12–14,18,19,21,24,25,29,35–37,39–42,44,45} (18 out of 35) reported estimates of associations adjusted for potential confounders.

Results of included studies

The characteristics of patients included in the studies and association of CCI score on outcomes are described in *Table 4*.

Acute coronary syndrome

A total of 11 studies^{11,17,18,21,22,28,32,33,36,37} evaluated the impact of co-morbidity in 1 154 408 patients admitted with ACS; however, only 5 studies^{11,21,28,31,37} reported on patients with no co-morbidity (37% of patients had CCI = 0). Five studies^{17,18,31,32,36} were statistically pooled for the association between an incremental increase in CCI and mortality (*Figure 3A*). Among patients with ACS, the risk of death was significantly greater with incremental increase in CCI score (RR 1.33; 95% CI 1.15–1.54). Three studies ($I^2 = 96\%$)^{11,21,31} compared patients with no co-morbidity (CCI score of 0) vs. patients with any co-morbidity (CCI score of >0) showing that the presence of co-morbidity (CCI score of >0) resulted in almost twice the risk of death (RR 1.93; 95% CI 1.67–2.24). Radovanovic *et al.*³¹ and Huang *et al.*²¹ also analysed the impact of CCI score of 0–1 vs. >1 showing a higher risk of death in patients



with CCI score of >1 (RR 2.26; 95% CI 1.23–4.16; $I^2 = 98\%$). Three studies^{21,31,37} demonstrated a more than two-fold rise in mortality in patients with a CCI score of >2 comparing with a score

of 0–2. Only one study¹⁶ compared CCI score of 0–3 vs. >3 , which reported higher mortality (RR 5.89; 95% CI 5.56–6.24) in patients with more co-morbidities (CCI score of >3).




Figure 2 Charlson co-morbidity individual component distribution.




Figure 2 Continued.

In an ACS registry (AMIS registry), Jeger *et al.*²² reported an increased risk of MACE (a composite endpoint of re-infarction, CVA, and/or death) over a 1-year follow-up period in patients with CCI score of ≥ 2 . In another study, Núñez *et al.*²⁸ demonstrated that a higher CCI score was an independent predictor of mortality or acute myocardial infarction at 30 days and 1 year.

Stable coronary heart disease

Two studies^{14,35} studied the relationship between incremental rise in CCI score and mortality in patients with stable CHD (Figure 3B), suggesting that incremental increases in CCI score were associated with worse outcomes (RR 1.38; 95% CI 1.29–1.48; $I^2 = 0\%$). Sachdev *et al.*³⁵ also reported that patients with a CCI score of 0 have better long-term survival (RR 1.88; 95% CI 1.48–2.38). They also reported that almost half of the patients (49%) included in the cohort were disease free and had no co-morbidities (CCI = 0).

Patients undergoing percutaneous coronary intervention


Lastly, five studies^{16,20,25,38,44} reported impact of CCI on long-term survival in patients undergoing PCI, out of which four indicated that mortality increases with each point rise in CCI score (RR 1.21; 95%

CI 1.12–1.31; $I^2 = 71\%$) (Figure 3C). Only Mamas *et al.*²⁵ reported about patients with no co-morbidities in their study.

Heart failure

A total of 13 studies reported the influence of co-morbidity in 63 609 patients with an underlying diagnosis of heart failure. An increased risk of mortality (RR 1.21; 95% CI 1.13–1.29; $I^2 = 48\%$) was observed per point increase in CCI score among four studies.^{13,15,26,41} Jong *et al.*²³ and Rodriguez-Pascual *et al.*³⁴ compared patients with CCI score of 0–1 vs. >1 and demonstrated that a CCI score of >1 was associated with an increased risk of death (RR 1.60; 95% CI 1.52–1.70; $I^2 = 0\%$). Similar trends were observed in studies that compared a CCI score of >2 with a CCI score of 0–2. For instance, three studies^{23,29,30} reported an increased risk of death (RR 1.76; 95% CI 1.65–1.87; $I^2 = 0\%$) in patients with CCI score of >2 . Patients with high burden of co-morbidities (CCI score of >4) were analysed in three studies,^{29,34,42} which showed almost three-fold increase in relative risk of mortality (RR 2.93; 95% CI 1.99–4.31; $I^2 = 15\%$). Two studies^{27,45} reported increased risk of death with higher co-morbid burden with hazard ratio of >1 , but it was unclear how they are related to CCI score. Both studies

A Acute coronary syndrome



B Stable angina/stable coronary heart disease




Figure 3 Acute coronary syndrome patients, stable angina/stable coronary heart disease patients, and patients undergoing percutaneous coronary intervention and mortality according to Charlson Co-morbidity Index. (A) Acute coronary syndrome. (B) Stable angina/stable coronary heart disease. (C) Patients undergoing PCI.

Table 3 Quality of included studies

Study ID	Prospective study design	Reliable ascertainment of outcomes	Less than 10% loss to follow-up	Use of adjustments for potential confounders
Bottle 2013 ¹¹	No; retrospective	Yes; death from death certificates from the Office for National Statistics	Unclear	None
Bar 2011 ¹²	Yes; prospective	Unclear; outcome assessed using modified Rankin scale out to 12 months by unclear methods	Unclear	Adjusted for presence of IVH, intratentorial ICH, and use of early DNACPR orders
Chin 1998 ¹³	Yes; prospective	Yes; death from chart review, survey of families, and search of the National Death Index	Yes; 7 patients discharged quickly and unreachable, 5 too sick for interview	White ethnicity, age ≥ 70 years, prior congestive heart failure, chronic pulmonary disease, Charlson Co-morbidity Index score, third heart sound, serum sodium ≤ 135 , EF < 0.50 , diabetes, respiratory rate $> 30/\text{min}$, cardiomegaly on admission chest radiograph
Chirinos 2006 ¹⁴	Yes; prospective	Yes; patients' interview and review of hospital electronic records	Yes; 9 patients were lost to follow-up	Multivariate analysis adjustments for age, left ventricular ejection fraction, congestive heart failure, and number of coronary artery territories involved with haemodynamically significant lesions
Clarke 2011 ¹⁵	No; retrospective	Yes; electronic database, review of medical notes, clinic visits, and review of death certificates	Yes; none	None
Eberti 2013 ¹⁶	Yes; prospective	Unclear; 1-year all-cause and cardiac mortality by unclear method	Unclear	None
Erickson 2014 ¹⁷	No; retrospective	Yes; 6-month post-discharge all-cause mortality or secondary cardiovascular events or revascularization procedures	Yes; none	None
Fabbian 2013 ¹⁸	No; retrospective	Yes; in-hospital mortality for myocardial infarction	Yes; none	Chronic kidney disease
Goldstein 2004 ¹⁹	Yes; prospective	Yes; death at discharge and 1-year mortality	Yes; none	Initial stroke severity
Hong 2011 ²⁰	Yes; prospective	Yes; followed up at 30, 180, and 360 days by telephone communication, office visit, or by contacts with primary physicians or referring cardiologists for 1-year mortality, stent thrombosis	Yes; none	Unclear
Huang 2015 ²¹	No; retrospective	Yes; data collected from National Health Insurance Research Database and the National Disability Registration Database of Taiwan	Yes; none	Adjusted (Model A); variables unclear
Jeger 2014 ²²	No; retrospective	Yes; data collected from AMIS plus registry	Yes; 161 lost to follow-up	None
Jong 2002 ²³	No; retrospective	Yes; 30-day and 1-year mortality ascertained by linking the database with Ontario registered person database	Yes; none	None
Khawaja 2014 ²⁴	Unclear	Unclear; primary outcomes of modified Rankin scale of 4–6, death and poor discharge disposition (any disposition other than home or inpatient rehabilitation) assessed by unclear methods	Unclear	Adjusted for baseline ICH score
Mamas 2015 ²⁵	Yes; prospective	Yes; data was collected into a Web-based data management system and an independent clinical events committee adjudicated all events	No; 326 lost to follow-up at 5 years	Adjusted for baseline demographic and lesion characteristic variables with a P -value of < 0.05
Menendez-Colino 2013 ²⁶	Unclear	Unclear; mortality at 12 months; unclear follow-up methods	Yes; 25 patients	Unclear
Munoz-Rivas 2009 ²⁷	No; retrospective	Unclear	Unclear	Unclear
Núñez 2004 ²⁸	Yes; prospective	Yes; 30-day and 1-year mortality or reinfarction at outpatient follow-up and telephonic contact	Yes; none	None
Oudejans 2012 ²⁹	Yes; prospective	Yes; all-cause mortality within 3 years; follow-up information obtained from hospital information system or from patient's general practitioners	Yes; 1 patient was lost to follow-up	Age, gender, LVEF, and NT-proBNP

Perez-Barquero 2010 ³⁰	No; retrospective	Unclear; in-hospital mortality by unclear follow-up methods	Unclear	Unclear	
Radovanovic 2014 ³¹	Yes; prospective	Yes; data collected from AMIS plus registry	No; 1091 patients' CCI data were not available	None	
Ramirez-Marrero 2011 ³²	No; retrospective	Unclear	Yes; none	None	
Ramirez-Marrero 2013 ³³	No; retrospective	Yes; cardiovascular mortality during follow-up	Yes; none	None	
Rodriguez-Pascual 2012 ³⁴	Yes; prospective	Unclear; mortality	Unclear	None	
Sachdev 2004 ³⁵	Yes; prospective	Yes; patients were followed up at 6 months, 1 year, and then annually by a mailed questionnaire, with telephone backup, as well as a National Death Index search for non-responders through December 2000	Yes; none	Adjusted for age, unclear if other variables were adjusted	
Sanchez 2011 ³⁶	Yes; prospective	Yes; data collected from admission records and follow-up	Yes; 4 patients did not complete follow-up	Adjusted for variables with $P < 0.05$ but variables unclear	
Schmidt 2012 ³⁷	No; retrospective	Yes; standardized incidence rate of myocardial infarction and 30-day and 31- to 365-day mortality by sex	Unclear	Age and sex	
Singh 2011 ³⁸	Yes; prospective	Yes; all-cause mortality during follow-up; the second main outcome was MI defined as the presence of 2 of 3 following criteria: prolonged (> 20 min) ischaemic chest pain and elevation of cardiac biomarkers (creatinine kinase-MB or relative index) more than two times upper limit of normal, or electrocardiographic changes (ST/T-wave changes or new Q waves)	Yes; 2% participants lost to follow-up	None	
Subramanian 2007 ³⁹	Yes; prospective	Yes; 5-year mortality during follow-up data obtained from Veterans Integrated Health Systems Technology Architecture databases	Yes; 35 patients were excluded for missing values	Adjusted; variables unclear	
Teng 2014 ⁴⁰	Yes; prospective	Yes; data were collected from the Hospital Morbidity Data Collection, which is linked to the Mortality register	Unclear	Adjusted; variables unclear	
Testa 2009 ⁴¹	Yes; prospective	Yes; all subjects were contacted at home or in their institution and examined by physicians trained to administer a questionnaire	Yes; 35 patients were unreachable and 9 did not have social support	Age, sex, heart rate, systolic blood pressure, diastolic blood pressure, social support, drugs number, MMSE, BADL, NYHA, CAD, chronic obstructive pulmonary disease, neurological disease, congestive heart failure, and CCI	
Theuns 2011 ⁴²	Yes; prospective	Yes; the data collected from two prospective ICD registries from Rotterdam and Basel; patient followed up at outpatient clinics	Yes; none	Adjusted for age	
Tuttolomondo 2008 ⁴³	Yes; prospective	Yes; demographic data and follow-up were collected from GIFA registry	Yes; none	None	
Urban 2011 ⁴⁴	Yes; prospective	Yes; the data collected from the e-Select registry where patients were followed up at 30, 180, and 360 days by telephone communication or office visit by contacts with primary physicians or referring cardiologist	Unclear	Adjusted for variables with entry P -value of 0.10 and stay criterion of 0.15; unclear exact variables	
Van Wijk 2013 ⁴⁵	Yes; prospective	Yes; clinically followed up for 18 months with recording of hospitalization, mortality, and adverse events up to 5 years	Unclear	Adjusted; variables unclear	

Table 4 Follow-up and results of the association between Charlson Co-morbidity Index and outcome

Study ID	Type of population (CAD, HF, CVA)	Definition of CCI	Outcome and duration of follow-up	Results demonstrating association between CCI and outcome
Bottle 2013 ¹²	ACS (CAD)	Charlson score 0 vs. >0	30-day mortality	30-day mortality: CCI score 0: 8370/151 577 (5.5%); CCI score >0: 20 999/177 792 (11.8%)
Bar 2011 ¹¹	Stroke (CVA)	Incremental rise in CCI from 0 to >3	12-month functional outcome according to modified Rankin scale	CCI score 1: OR 1.78 (0.86–3.70); CCI score 2: OR 2.34 (0.98–5.61); CCI score 3: OR 3.48 (1.64–7.37)
Chin 1998 ¹³	HF	Incremental increase in CCI	Time to mortality	Mortality per CCI point to max of 4 points: HR 1.3 (1.1–1.4)
Chirinos 2006 ¹⁴	Stable CAD	Incremental increase in modified CCI	All-cause mortality during 58-month follow-up	Odds of mortality with incremental increase in modified CCI score: OR 1.32 (1.17–1.48)
Clarke 2011 ¹⁵	Heart failure (HF)	Incremental increase in CCI	Time to mortality with follow-up of mean of 4.4 years	Overall mortality by per unit increase in CCI: HR 1.26 (1.19–1.35)
Eberti 2013 ¹⁶	PCI (CAD)	Mortality by different CCI score	1-year mortality and cardiac mortality	Overall 1-year mortality: CCI score 0: 182041 (0.9%); CCI score 1: 282162 (1.3%); CCI score 2: 187776 (2.3%); CCI score ≥3: 251578 (4.3%)
Erickson 2014 ¹⁷	ACS (CAD)	Incremental increase in CCI	Inpatient and 6-month mortality and post-discharge cardiac event or procedure	Cardiac mortality: CCI score 0: 147041 (0.7%); CCI score 1: 1372162 (0.6%); CCI score 2: 91776 (1.2%); CCI score ≥3: 141578 (2.4%)
Fabbian 2013 ¹⁸	ACS (CAD)	Incremental increase in CCI	In-hospital mortality from MI	Inpatient death with CCI: OR 1.28 (1.14–1.43) 6-month death with CCI: OR 1.55 (1.41–1.72) Post-discharge cardiac event or procedure CCI: 1.21 (1.12–1.31) In-hospital mortality for MI with CCI without renal dysfunction: OR 1.101 (1.069–1.134)
Goldstein 2004 ¹⁹	Stroke (CVA)	Low CCI 0–1 vs. high CCI ≥2	1-year mortality	1-year mortality with low CCI score 0–1: 88/551 (16%) High CCI score ≥2: 106/429 (26%)
Hong 2011 ²⁰	PCI (CAD)	Incremental rise in CCI on outcomes	Time to mortality or stent thrombosis with follow-up up to 1-year	Every 1-point increment in CCI on death: HR 1.3 (1.1–1.5) Every 1-point increment on stent thrombosis: HR 1.5 (1.3–1.8)
Huang 2015 ²¹	ACS (CAD)	Risk for each CCI score	Time to acute myocardial infarction	Adjusted model A (unclear variables): CCI score 1: HR 2.25 (2.12–2.39); CCI score 2: HR 3.07 (2.89–3.26); CCI score 3: HR 3.71 (3.48–3.95); CCI score ≥4: HR 5.89 (5.56–6.25)
Jeger 2014 ²²	ACS (CAD)	Charlson score ≥2	1-year MACE	1-year MACE with CCI score ≥2: OR 1.42 (1.05–1.92)
Jong 2002 ²³	Heart failure (HF)	CCI score and mortality rate	30-day and 1-year mortality	CCI score 0: 30-day mortality 1397/15 020 (9.3%); 1-year mortality 4025/15 020 (26.8%) CCI score 1: 30-day mortality 1348/12 602 (10.7%); 1-year mortality 3907/12 602 (31.0%)
Khawaja 2014 ²⁴	Stroke (CVA)	Incremental increase in CCI score	Death at unclear follow-up	CCI score 2: 30-day mortality 895/6485 (13.8%); 1-year mortality 2555/6485 (39.4%) CCI score 3: 30-day mortality 864/4595 (18.8%); 1-year mortality 2325/4595 (50.6%) Death and CCI score: OR 1.05 (0.91–1.21)
Mamas 2015 ²⁵	PCI (CAD)	Incremental increase in CCI score	30-day, 1-year, and 5-year cardiac death and MACE	30-day: cardiac death: OR 1.47 (1.20–1.80); MACE: OR 1.27 (1.11–1.44) 1-year: cardiac death: OR 1.46 (1.30–1.65); MACE: OR 1.32 (1.23–1.42) 5-year: cardiac death: OR 1.38 (1.24–1.53); MACE: OR 1.29 (1.22–1.36)
Menendez-Colino 2013 ²⁶	Heart failure (HF)	CCI score and mortality	Time to mortality with follow-up maximum of 12 months	CCI score: HR 1.13 (1.04–1.24)
Munoz-Rivas 2009 ²⁷	Heart failure (HF)	Incremental increase in CCI	Survival	Survival with incremental CCI: HR 1.46 (1.21–5.07)

Núñez 2004 ²⁸	ACS (CAD)	CCI score and risk compared with CCI score 0	Time to death or reinfarction to a maximum of 30 days and 1 year	Risk of death or reinfarction at 30 days: CCI score 1: HR 1.00; CCI score 2: HR 1.69 (1.10–2.59); CCI score 3: HR 1.78 (1.08–2.92); CCI score 4: HR 1.57 (0.87–2.83) Risk of death or reinfarction at 1 year: CCI score 1: HR 1.00; CCI score 2: HR 1.62 (1.18–2.23); CCI score 3: HR 2.00 (1.39–2.89); CCI score 4: HR 2.24 (1.50–3.36) 3-year mortality: CCI score 0–2: HR 1.00; CCI score 3–4: HR 1.5 (0.7–2.9); CCI score >4: HR 4.0 (1.9–8.8)
Oudejans 2012 ²⁹	Heart failure (HF)	CCI score 0–2 vs. 3–4 or ≥4	Time to mortality to a maximum of 3 years	In-hospital mortality: CCI score 1–2: 76/1528; CCI score ≥3: 48/599
Perez-Barquero 2010 ³⁰	Heart failure (HF)	CCI score 1–2 vs. ≥3	In-hospital mortality	In-hospital mortality compared with CCI score 0: CCI score 1: OR 1.36 (1.16–1.60); CCI score 2: OR 1.65 (1.38–1.97); CCI score ≥3: OR 2.20 (1.86–2.57)
Radovanovic 2014 ³¹	ACS (CAD)	Incremental rise in CCI and risk compared with CCI score 0	In-hospital mortality and 1-year mortality assessed using data from AMIS plus registry	1-year mortality per CCI point: age-adjusted mortality: OR 1.44 (1.36–1.53)
Ramirez-Marrero 2011 ³²	ACS (CAD)	Higher CCI treated as incremental	In-hospital mortality and median follow-up of 24 months	In-hospital mortality: OR 1.6 (1.4–1.8); long-term mortality: OR 1.3 (1.2–1.5); readmission for HF: OR 1.2 (1.04–1.3); MACE during follow-up: OR 1.1 (1–1.2)
Ramirez-Marrero 2013 ³³	ACS (CAD)	Highest CCI score	Cardiovascular mortality during follow-up of 36 months	CCI and long-term mortality: OR 1.72 (1.09–2.71)
Rodriguez-Pascual 2012 ³⁴	Heart failure (HF)	CCI score	Mortality	Mortality by CCI score: 0–1: 5/121; 2–4: 17/227; ≥5: 26/194
Sachdev 2004 ³⁵	Stable CAD	CCI scores of 0, 1, and ≥2	Time to mortality during follow-up period of almost 11 years	CCI score 0: 95/810 (11.7%); CCI score 1: 58/378 (15.3%); CCI score ≥2: 88/283 (31.1%) Incremental increase in modified CCI: HR 1.41 (1.30–1.53) Per point increase in CCI: HR 1.3 (1.2–1.4)
Sanchis 2011 ³⁶	ACS (CAD)	Incremental increase in CCI per point	Time to mortality to a maximum of 1 year	30-day mortality: RR 1.96 (1.83–2.11) 31- to 365-day mortality: RR 3.89 (3.58–4.24)
Schmidt 2012 ³⁷	ACS (CAD)	CCI score 0 (normal) vs. ≥3 (very severe)	Time to mortality or myocardial infarction during median follow-up of 35 months	Death during follow-up: HR 1.12 (1.06–1.18)
Singh 2011 ³⁸	PCI (CAD)	Incremental increase in CCI per point	Time to mortality at follow-up of up to 5 years	Death/MI during follow-up: HR 1.05 (1.01–1.10) 5-year all-cause mortality: HR 1.39 (1.16–1.67)
Subramanian 2007 ³⁹	Heart failure (HF)	Incremental increase in 3 points of CCI	Time to mortality at follow-up of up to 5 years	1-year mortality with CCI: <55 years: HR 1.38 (1.26–1.51); ≥55 years: HR 1.20 (1.18–1.22)
Teng 2014 ⁴⁰	Heart failure (HF)	CCI unclear if incremental or cut-off	1-year mortality	12-year mortality with CCI: HR 1.15 (1.01–1.31)
Testa 2009 ⁴¹	Heart failure (HF)	Incremental increase in CCI score	Time to mortality to a maximum follow-up of 12 years	All-cause mortality: HR 3.49 (2.06–6.60)
Theuns 2011 ⁴²	Heart failure (HF)	CCI score >5	Time to all-cause mortality during a median follow-up of 30.5 months	In-hospital mortality: OR 35.7 (4.8–265.2)
Tuttolomondo 2008 ⁴³	Stroke (CVA)	CCI score <2 vs. CCI score >2	In-hospital mortality	1-year death: HR 1.2 (1.1–1.2) 1-year stent thrombosis: HR 1.2 (1.1–1.4) 1-year major bleeding: HR 1.1 (1.0–1.2)
Urban 2011 ⁴⁴	PCI (CAD)	Incremental increase in CCI per point	Time to death, stent thrombosis, and major bleeding at maximum of 1 year	CCI score: HR 2.47 (1.27–4.83)
Van Wijk 2013 ⁴⁵	Heart failure (HF)	Incremental increase in CCI score	Hospital-free survivals during follow-up period	

C Patients undergoing PCI




Figure 3 Continued.

were only available in abstract form and, therefore, not included in the final meta-analysis. More interestingly, Subramanian et al.³⁹ assessed the impact of incremental increase in CCI per 3 points in heart failure patients over 5 years, reporting increased risk of death (HR 1.39; 95% CI 1.16–1.67) with growing burden of co-morbidities.

Cerebrovascular accident

A total of four studies analysed the impact of CCI score on survival in patients with an acute CVA. Khawaja et al.²⁴ reported a no significant increased risk of death with incremental increase in CCI score (RR 1.05; 95% CI 0.91–1.21). However, higher CCI score (>2) had significant impact on mortality (RR 3.80; 95% CI 1.20–12.01; $I^2 = 84\%$) when compared with low CCI score (0–2).

Discussion

In this study, we evaluated the prevalence and prognostic impact of co-morbidities as defined by CCI in patients with CHD, heart failure, and CVA. We observed a significant burden of co-morbidity in patients with CVD—two-thirds of patients included in the analysis had at least one chronic condition. The most common cardiovascular co-morbid conditions identified in patients with CHD were diabetes and history of prior myocardial infarction, whereas chronic obstructive pulmonary disease and kidney disease were the most frequent non-cardiovascular conditions. To our knowledge, this is the first study to systematically show the impact of co-morbid burden as defined by CCI on survival in patients with CHD, heart

failure, and CVA. We found that the presence of co-morbidities had a significant incremental prognostic impact in patients with a broad range of CVDs.

CHD is the commonest CVD affecting one in seven people in USA⁴⁶ and UK every year. Patients with CHD are likely to have higher number of coexisting illnesses either in the form of prevalent cardiovascular risk factors such as diabetes and hypertension or in the form of direct manifestations of CHD such as prior myocardial infarction or heart failure. For instance, in one study, diabetes, hypertension, and heart failure were found to be most frequently encountered coexisting illnesses in patients admitted with ACS and 68% of the participants had at least three co-morbidities.⁴⁷ The rising burden of co-morbidity has been reported to have inverse relationship with survival outcomes in patients with CHD. In our analysis, incremental rise in CCI was associated with significant increase in mortality and the risk of death was almost doubled with the presence of any co-morbidity compared with the patients with no co-morbidity (Figure 3A). This has important clinical implications in this cohort of patients as the prevalent cardiovascular risk factors such as hyperlipidaemia, smoking, and other related cardiovascular co-morbidities such as hypertension and diabetes in patients with CHD are usually treated aggressively, but there is growing evidence that non-CVD burden may also contribute to increased risk of mortality.^{25,31} We report that CCI is not only a simple way of quantifying co-morbid burden but also provides prognostic value in ascertaining outcomes. Clinicians often use risk assessment tools such as GRACE and TIMI scores in determining

the type of intervention, treatment plan, and allocation of resources in managing patients with ACS. Although these models have been validated in predicting the adverse events,^{48,49} the clinical data incorporated in these models do not take into account the co-morbid burden of the patients. Previous studies have suggested that the performance of such risk models improves when co-morbidity scores such as CCI are added to the risk scores¹⁷ and may help in better allocations of resources and developing robust treatment pathways for patients with multiple co-morbidities. Our study highlights the importance of taking into consideration of the overall co-morbid burden in such patients while making the therapeutic decisions. Furthermore, our study also demonstrates that co-morbidity burden has prognostic value.

The prevalence of heart failure is increasing due to the ageing population and better survival from acute cardiac events.⁵⁰ Our findings reinforce the hypothesis that heart failure patients with multiple co-morbidities have worse outcomes.¹⁵ Similarly, increasing co-morbid burden is associated with a worse prognosis in patients after an acute cerebrovascular event. We observed that the risk of death was almost four-fold greater in patients with two or more coexisting illnesses (Figures 4 and 5).

The mechanism by which the coexisting co-morbid burden influences outcomes in patients with CVD is complex and multifactorial. Older and frailer patients with high burden of co-morbidities are more likely to be treated conservatively following a cardiovascular event.^{51,52} For instance, a large national ACS registry reported an

incremental reduction in provision of evidence-based treatments such as aspirin, statins, angiotensin-converting enzyme (ACE) inhibitors, and reperfusion therapy to the older multi-morbid patients.⁵³ In another recent analysis of 18 814 patients, Patel *et al.* identified that patients with higher co-morbid burden as defined by CCI were less likely to receive coronary artery angiography and/or revascularization following presentation with ST-segment elevation myocardial infarction.⁵⁴ Similarly, thrombolysis therapy in acute ischaemic stroke is usually reserved for younger patients with no significant burden of co-morbidities due to fear of less favourable outcomes such as bleeding complications in elderly patients with multiple co-morbidities.⁵⁵ In the management of patients with chronic heart failure, the associated burden of co-morbidities may limit the use of medications such as ACE inhibitors or spironolactone, particularly in patients with severe chronic kidney disease⁵⁶ and β -blockers in patients with coexisting severe chronic obstructive pulmonary disease. Furthermore, patients with multiple chronic conditions are less likely to receive invasive therapies such as implantable cardioverter-defibrillators or cardiac resynchronization therapy.⁵⁷ There is also growing evidence that increasing burden of co-morbidities in patients with heart failure is associated with repeated hospitalization and poor outcomes.^{58,59}

Provision of aggressive treatment strategies in patients with multi-morbidity can lead to higher incidence of complications and adverse outcomes. For example, patients with leukaemia are at higher risk of stent thrombosis⁶⁰ and those with liver dysfunction are at increased




Figure 4 Heart failure patients and mortality according to Charlson Co-morbidity Index.




Figure 5 Cerebrovascular accident patients and mortality according to Charlson Co-morbidity Index.

risk of bleeding complications post-PCI and cardiac mortality.⁶¹ Similarly, the presence of diabetes and haematological disorders has been shown to increase the risk of haemorrhagic transformation in patients with ischaemic stroke.^{62,63} Consequently, the presence of coexisting diseases may drive poor outcomes in patients with CHD due to reduced scope of treatment options and increased risk of complications. Hence, clinicians may be reserved in deciding treatment strategies while managing patients with multi-morbidity due to the challenge of finding a balance between risk and benefit of an intervention.^{64,65}

Other factors that may be responsible for deleterious effect of co-morbidities on survival outcomes are the presence of coexisting illness sharing the same pathophysiology and adverse drug reactions due to polypharmacy. For example, the presence of anaemia results in low cardiac output state and has been reported to have synergetic impact on the mortality in patients with chronic heart failure.⁶⁶

Our findings have important implications in the management of patients with CHD, heart failure, and cerebrovascular disease. Treatment options such as medical therapies, PCI, surgical revascularization, device therapies, and thrombolysis are now readily available to wider spectrum of patients. Although international guidelines^{67,68} advocate a comprehensive assessment of patients taking into account their co-morbid status, contemporary risk stratification tools such as GRACE, Cath PCI, and Syntax are derived from data sets based on patient's characteristics, procedural demographics, and cardiovascular risk factors and do not take into account patients' co-morbid burdens. Our analysis shows that CCI score has prognostic value in our cohort of patients and using CCI alongside these risk models can help physicians to ascertain outcomes and better resource allocation. For instance, the addition of CCI to the Mayo Clinic Risk Score for PCI increased net reclassification index by 34% and improved the c-statistic for the model significantly.³⁸ Erickson et al.¹⁷ also tested the risk prediction of GRACE model by adding CCI and observed a significant improvement in predicting outcomes in ACS patients. Another study reported improved discriminative performance of GRACE Risk Prediction Index score when added with CCI in predicting future cardiac-related events post-myocardial infarction.¹⁷ Therefore, the

assessment of co-morbid status and its impact on long-term survival should be integrated into the counselling of the patients before deciding the choice of treatment in conjunction with traditional risk assessment.

Our study has several strengths and limitations. To our knowledge, this is the first review on impact of co-morbidity defined by CCI on major CVD such as CHD, heart failure, and cerebrovascular disease. We were able to analyse the impact of per unit rise in CCI in our cohort of patients demonstrating that rise in CCI score has inverse relationship with survival. We were also able to evaluate the impact of CCI among individual cohorts of CHD namely stable angina, ACS, and those undergoing PCI and found a uniform negative impact of rising CCI score across all cohorts. Additionally, we also studied the prevalence of co-morbidities in patients with CVD and found that majority of patients in this cohort have significant burden of co-morbidities.

Our study was limited by the incomplete reporting of original studies and was reliant on the published data available. We were not able to evaluate the impact of individual components of CCI on mortality, as this was not consistently reported across all studies. Furthermore, the studies included in our review were mainly observational, which have their own inherent limitations and may be subject to selection biases and unmeasured confounders. Another limitation is that we found significant heterogeneity in several analyses. This may be because many of the studies are large with very narrow confidence intervals leading to statistical heterogeneity when there is little overlap in 95% CI among the studies. However, all the studies, in general, report estimates that are consistently significant and favour increased events with higher CCI score. The statistical heterogeneity arises from differences in each study in terms of population evaluated and study methodology, which leads to variation in estimates for the prognostic value of CCI.

Conclusion

Our study shows that co-morbid burden defined by CCI is significant across a broad range of cardiovascular conditions and has significant impact on survival in patients with CHD, heart failure,

and CVA. Assessment of co-morbid burden using CCI provides a method of quantifying risk associated with co-morbidities in patients with CVD and should be incorporated into decision-making processes when counselling patients regarding risk and benefits of treatment in conjunction with allocation of resources.

Supplementary material

Supplementary material is available at *European Heart Journal – Quality of Care and Clinical Outcomes* online.

Conflict of interest: none declared.

References

- Mendis S, Puska P, Norrving B, eds. *Global Atlas on Cardiovascular Disease Prevention and Control: Policies, Strategies and Interventions*. Geneva: World Health Organization in collaboration with World Heart Federation and World Stroke Organization; 2011. http://www.who.int/cardiovascular_diseases/publications/atlas_cvd/en/.
- Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;**380**:37–43.
- Wolff JL, Starfield B, Anderson G. Prevalence, expenditures, and complications of multiple chronic conditions in the elderly. *Arch Intern Med* 2002;**162**:2269–2276.
- Alexander KP, Newby LK, Cannon CP, Armstrong PW, Gibler WB, Rich MW, Van de Werf F, White HD, Weaver WD, Naylor MD, Gore JM, Krumholz HM, Ohman EM. Acute coronary care in the elderly, Part I: non-ST-segment-elevation acute coronary syndromes: a scientific statement for healthcare professionals from the American Heart Association Council on Clinical Cardiology: in collaboration with the Society of Geriatric Cardiology. *Circulation* 2007;**115**:2549–2569.
- Alexander KP, Newby LK, Armstrong PW, Cannon CP, Gibler WB, Rich MW, Van der Werf F, White HD, Weaver WD, Naylor MD, Gore JM, Krumholz HM, Ohman EM. Acute coronary care in the elderly, Part II: ST-segment-elevation myocardial infarction: a scientific statement for healthcare professionals from the American Heart Association Council on Clinical Cardiology: in collaboration with the Society of Geriatric Cardiology. *Circulation* 2007;**115**:2570–2589.
- McManus DD, Nguyen HL, Saczynski JS, Tisminetzky M, Bourell P, Goldberg RJ. Multiple cardiovascular comorbidities and acute myocardial infarction: temporal trends (1990–2007) and impact on death rates at 30 days and 1 year. *Clin Epidemiol* 2012;**4**:115–123.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;**40**:373–383.
- Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol* 1992;**45**:613–619.
- Deeks JJ, Higgins JPT, Altman DG, eds. Chapter 9: Analysing data and undertaking meta-analyses. In: Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0 (updated March 2011). The Cochrane Collaboration; 2011. Available from www.cochrane-handbook.org.
- Ioannidis JP, Trikalinos TA. The appropriateness of asymmetry tests for publication bias in meta-analyses: a large survey. *CMAJ* 2007;**176**:1091–1096.
- Bottle A, Sanders RD, Mozid A, Aylin P. Provider profiling models for acute coronary syndrome mortality using administrative data. *Int J Cardiol* 2013;**168**:338–343.
- Bar B, Hemphill JC. Charlson comorbidity index adjustment in intracerebral hemorrhage. *Stroke* 2011;**42**:2944–2946.
- Chin MH, Goldman L. Gender differences in 1-year survival and quality of life among patients admitted with congestive heart failure. *Med Care* 1998;**36**:1033–1046.
- Chirinos JA, Veerani A, Zambrano JP, Schob A, Perez G, Mendez AJ, Chakko S. Evaluation of comorbidity scores to predict all-cause mortality in patients with established coronary artery disease. *Int J Cardiol* 2007;**117**:97–102.
- Clarke B, Howlett J, Sapp J, Andreou P, Parkash R. The effect of comorbidity on the competing risk of sudden and nonsudden death in an ambulatory heart failure population. *Can J Cardiol* 2011;**27**:254–261.
- Eberli F, Valdes M, Menown I, Alhaddad I, Oldroyd K, Roffi M, Hildick-Smith D, Iosseliani D, Kapur K, Urban P. Comorbidities determine prognosis in patients undergoing coronary stenting: results from two large registries evaluating biolimus-A9-eluting from biodegradable polymer stents. *Eurointervention* 2013;**9**:194.
- Erickson SR, Cole E, Kline-Rogers E, Eagle KA. The addition of the Charlson Comorbidity Index to the GRACE Risk Prediction Index improves prediction of outcomes in acute coronary syndrome. *Popul Health Manag* 2014;**17**:54–59.
- Fabbian F, Pala M, De Giorgi A, Manfredini F, Menegatti AM, Salmi R, Portaluppi F, Gallerani M, Manfredini R. In-hospital mortality in patients with renal dysfunction admitted for myocardial infarction: the Emilia-Romagna region of Italy database of hospital admissions. *Int Urol Nephrol* 2013;**45**:769–775.
- Goldstein LB, Samsa GP, Matchar DB, Horner RD. Charlson Index comorbidity adjustment for ischemic stroke outcome studies. *Stroke* 2004;**35**:1941–1945.
- Hong YJ, Jeong MH, Abizaid A, Banning A, Bartorelli A, Dzavik V, Ellis SG, Gao R, Holmes DR Jr, Legrand V, Neumann FJ, Spaulding C, Worthley S, Urban P. Sirolimus-eluting coronary stents in octogenarians: a 1-year analysis of the worldwide e-SELECT Registry. *JACC Cardiovasc Interv* 2011;**4**:982–991.
- Huang Y-Y, Kung P-T, Chiu L-T, Tsai W-C. Related factors and incidence risk of acute myocardial infarction among the people with disability: a national population-based study. *Res Dev Disabil* 2015;**36**:366–375.
- Jeger R, Jaguszewski M, Nallamothu BN, Lüscher TF, Urban P, Pedrazzini GB, Erne P, Radovanovic D. Acute multivessel revascularization improves 1-year outcome in ST-elevation myocardial infarction: a nationwide study cohort from the AMIS Plus registry. *Int J Cardiol* 2014;**172**:76–81.
- Jong P, Vowinkel E, Liu PP, Gong Y, Tu JV. Prognosis and determinants of survival in patients newly hospitalized for heart failure: a population-based study. *Arch Intern Med* 2002;**162**:1689–1694.
- Khawaja A, Boehme AK, Albright K, Kumar G, Arora K, Lysterly M, Harrigan M. The utility of Charlson Comorbidity Index in predicting short-term outcomes in patients with primary intracerebral hemorrhage. *Stroke* 2015;**46**:ATP326.
- Mamas MA, Fath-Ordoubadi F, Danzi GB, Spaepen E, Kwok CS, Buchan I, Peek N, de Belder MA, Ludman PF, Paunovic D, Urban P. Prevalence and impact of comorbidity burden as defined by the Charlson co-morbidity index on 30-day and 1- and 5-year outcomes after coronary stent implantation (from the Nobori-2 study). *Am J Cardiol* 2015;**116**:364–371.
- Menendez-Colino R, Condorhuanan P, Diez-Sebastian J, Mauleon-Ladrero C, Paredes E, Ferrero-Martinez A, Rodriguez-Pascual C. One-year mortality related factors in older-old patients admitted with heart failure in six Spanish hospitals. *Eur Geriatr Med* 2013;**4**:S32.
- Munoz Rivas N, Mendez M, Ortiz Alonso J, Nieto S, Audibert Mena L, Renieblas P. The role of comorbidities in elderly patients with heart failure: Implication on survival. *Eur J Heart Fail Suppl* 2009;**8**:Abstract 1106.
- Núñez JE, Núñez E, Fácila L, Bertomeu V, Llacer À, Bodí V, Sanchis J, Sanjuan R, Blasco ML, Consuegra L, Martinez A, Chorro FJ. Prognostic value of Charlson comorbidity index at 30 days and 1 year after acute myocardial infarction. *Rev Esp Cardiol* 2004;**57**:842–849.
- Oudejans I, Mosterd A, Zuithoff NP, Hoes AW. Comorbidity drives mortality in newly diagnosed heart failure: a study among geriatric outpatients. *J Card Fail* 2012;**18**:47–52.
- Montero P-BM, Conthe GP, Román SP, García AJ, Forteza-Rey J. Morbidity from patients admitted for heart failure in the departments of internal medicine. *Rev Clin Esp* 2010;**210**:149–158.
- Radovanovic D, Seifert B, Urban P, Eberli FR, Rickli H, Bertel O, Puhan MA, Erne P. Validity of Charlson Comorbidity Index in patients hospitalised with acute coronary syndrome. Insights from the nationwide AMIS Plus registry 2002–2012. *Heart* 2014;**100**:288–294.
- Ramirez-Marrero MA, Jimenez-Navarro M, De Teresa-Galvan E, De Mora-Martin M. The importance of the Charlson index in risk stratification in patients admitted for acute coronary syndrome without ST-segment elevation. *Eur J Cardiovasc Prev Rehabil* 2011;**18**(Suppl. 1):S113.
- Ramirez-Marrero M, Roldan-Jimenez M, Delgado-Prieto J, Jimenez-Navarro M, de Mora-Martin M. Reperfusion of complete percutaneous coronary revascularization on prognosis of elderly patients. *Intensive Care Med* 2013;**39**:S247.
- Rodríguez-Pascual C, Vilches-Moraga A, Paredes-Galán E, Ferrero-Marinez AI, Torrente-Carballido M, Rodríguez-Artalejo F. Comprehensive geriatric assessment and hospital mortality among older adults with decompensated heart failure. *Am Heart J* 2012;**164**:756–762.
- Sachdev M, Sun JL, Tsatis AA, Nelson CL, Mark DB, Jollis JG. The prognostic importance of comorbidity for mortality in patients with stable coronary artery disease. *J Am Coll Cardiol* 2004;**43**:576–582.
- Sanchis J, Núñez J, Bodí V, Núñez E, García-Alvarez A, Bonanad C, Regueiro A, Bosch X, Heras M, Sala J, Bielsa O, Llacer A. Influence of comorbid conditions on one-year outcomes in non-ST-segment elevation acute coronary syndrome. *Mayo Clin Proc* 2011;**86**:291–296.
- Schmidt M, Jacobsen JB, Lash TL, Bøtker HE, Sørensen HT. 25 year trends in first time hospitalisation for acute myocardial infarction, subsequent short and long term mortality, and the prognostic impact of sex and comorbidity: a Danish nationwide cohort study. *BMJ* 2012;**344**:e356.
- Singh M, Rihal CS, Lennon RJ, Spertus JA, Nair KS, Roger VL. Influence of frailty and health status on outcomes in patients with coronary disease undergoing percutaneous revascularization. *Circ Cardiovasc Qual Outcomes* 2011;**4**:496–502.

39. Subramanian U, Eckert G, Yeung A, Tierney WM. A single health status question had important prognostic value among outpatients with chronic heart failure. *J Clin Epidemiol* 2007;**60**:803–811.
40. Teng T-HK, Katzenellenbogen JM, Thompson SC, Sanfilippo FM, Knuiman M, Geelhoed E, Hobbs M, Bessarab D, Hung J. Incidence of first heart failure hospitalisation and mortality in Aboriginal and non-Aboriginal patients in Western Australia, 2000–2009. *Int J Cardiol* 2014;**173**:110–117.
41. Testa G, Cacciatore F, Galizia G, Della-Morte D, Mazzella F, Russo S, Ferrara N, Rengo F, Abete P. Charlson Comorbidity Index does not predict long-term mortality in elderly subjects with chronic heart failure. *Age Ageing* 2009;**38**:734–740.
42. Theuns DA, Schaer BA, Soliman OI, Altmann D, Sticherling C, Geleijnse ML, Osswald S, Jordaens L. The prognosis of implantable defibrillator patients treated with cardiac resynchronization therapy: comorbidity burden as predictor of mortality. *Europace* 2011;**13**:62–69.
43. Tuttolomondo A, Pedone C, Pinto A, Di Raimondo D, Fernandez P, Di Sciacca R, Licata G. Predictors of outcome in acute ischemic cerebrovascular syndromes: the GIFA study. *Int J Cardiol* 2008;**125**:391–396.
44. Urban P, Abizaid A, Banning A, Bartorelli AL, Baux AC, Džavík V, Ellis S, Gao R, Holmes D, Jeong MH, Legrand V, Neumann FJ, Nyakern M, Spaulding C, Worthley S. Stent thrombosis and bleeding complications after implantation of sirolimus-eluting coronary stents in an unselected worldwide population: a report from the e-SELECT (Multi-Center Post-Market Surveillance) registry. *J Am Coll Cardiol* 2011;**57**:1445–1454.
45. Van Wijk S, Bektas S, Muzzarelli S, Kiencke S, Maeder M, Estlinbaum W, Tobler D, Erne P, Pfisterer ME, Brunner-La R. Impact of comorbidities on safety, tolerability and efficacy of intensified medical therapy in heart failure. *Eur Heart J* 2013;**34**(Suppl. 1):P2742.
46. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, de Ferranti S, Després JP, Fullerton HJ, Howard VJ, Huffman MD, Judd SE, Kissela BM, Lackland DT, Lichtman JH, Lisabeth LD, Liu S, Mackey RH, Matchar DB, McGuire DK, Mohler ER 3rd, Moy CS, Muntner P, Mussolino ME, Nasir K, Neumar RW, Nichol G, Palaniappan L, Pandey DK, Reeves MJ, Rodriguez CJ, Sorlie PD, Stein J, Towfighi A, Turan TN, Virani SS, Willey JZ, Woo D, Yeh RW, Turner MB. Heart disease and stroke statistics—2015 update: a report from the American Heart Association. *Circulation* 2015;**131**:e29–322.
47. Taneva E, Bogdanova V, Shtereva N. Acute coronary syndrome, comorbidity, and mortality in geriatric patients. *Ann NY Acad Sci* 2004;**1019**:106–110.
48. Pollack CV, Sites FD, Shofer FS, Sease KL, Hollander JE. Application of the TIMI risk score for unstable angina and non-ST elevation acute coronary syndrome to an unselected emergency department chest pain population. *Acad Emerg Med* 2006;**13**:13–18.
49. Elbarouni B, Goodman SG, Yan RT, Welsh RC, Kornder JM, DeYoung JP, Wong GC, Rose B, Grondin FR, Gallo R, Tan M, Casanova A, Eagle KA, Yan AT. Validation of the Global Registry of Acute Coronary Event (GRACE) risk score for in-hospital mortality in patients with acute coronary syndrome in Canada. *Am Heart J* 2009;**158**:392–399.
50. Mosterd A, Hoes AW. Clinical epidemiology of heart failure. *Heart* 2007;**93**:1137–1146.
51. Lee JH, Yang DH, Park HS, Cho Y, Jeong MH, Kim YJ, Kim KS, Hur SH, Seong IW, Hong TJ, Cho MC, Kim CJ, Jun JE, Park WH, Chae SC. Suboptimal use of evidence-based medical therapy in patients with acute myocardial infarction from the Korea Acute Myocardial Infarction Registry: prescription rate, predictors, and prognostic value. *Am Heart J* 2010;**159**:1012–1019.
52. Passman R, Kimmel S. Do comorbidities influence the treatment of myocardial infarction? *J Gen Intern Med* 1997;**12**:73.
53. Zaman MJ, Stirling S, Shepstone L, Ryding A, Flather M, Bachmann M, Myint PK. The association between older age and receipt of care and outcomes in patients with acute coronary syndromes: a cohort study of the Myocardial Ischaemia National Audit Project (MINAP). *Eur Heart J* 2014;**35**:1551–1558.
54. Patel N, Patel N, Thakkar B, Singh V, Arora S, Savani C, Deshmukh A, Thadani U, Badheka AO, Alfonso C, Fonarow GC, Cohen MG. Management strategies and outcomes of ST-segment elevation myocardial infarction patients transferred after receiving fibrinolytic therapy in the United States. *Clin Cardiol* 2016;**39**:9–18.
55. Bateman BT, Schumacher HC, Boden-Albala B, Berman MF, Mohr J, Sacco RL, Pile-Spellman J. Factors associated with in-hospital mortality after administration of thrombolysis in acute ischemic stroke patients: an analysis of the nationwide inpatient sample 1999 to 2002. *Stroke* 2006;**37**:440–446.
56. Muzzarelli S, Maeder MT, Toggweiler S, Rickli H, Nietlispach F, Julius B, Burkard T, Pfisterer ME, Brunner-La Rocca HP. Frequency and predictors of hyperkalemia in patients ≥60 years of age with heart failure undergoing intense medical therapy. *Am J Cardiol* 2012;**109**:693–698.
57. Green AR, Leff B, Wang Y, Spatz ES, Masoudi FA, Peterson PN, Daugherty SL, Matlock DD. Geriatric conditions in patients undergoing defibrillator implantation for prevention of sudden cardiac death prevalence and impact on mortality. *Circ Cardiovasc Qual Outcomes* 2016;**9**:23–30.
58. Braunstein JB, Anderson GF, Gerstenblith G, Weller W, Niefeld M, Herbert R, Wu AW. Noncardiac comorbidity increases preventable hospitalizations and mortality among Medicare beneficiaries with chronic heart failure. *J Am Coll Cardiol* 2003;**42**:1226–1233.
59. Muzzarelli S, Leibundgut G, Maeder MT, Rickli H, Handschin R, Gutmann M, Jeker U, Buser P, Pfisterer M, Brunner-La Rocca HP. Predictors of early readmission or death in elderly patients with heart failure. *Am Heart J* 2010;**160**:308–314.
60. Sargyan Z, Higgins C, Alexandrescu S, Ott DA, Jain SK. Acute promyelocytic leukemia as a cause of intracoronary drug-eluting–stent thrombosis. *Tex Heart Inst J* 2012;**39**:416.
61. Tanaka S, Sakata R, Marui A, Furukawa Y, Kita T, Kimura T. Predicting long-term mortality after first coronary revascularization—the Kyoto model. *Circ J* 2012;**76**:328–334.
62. Paciaroni M, Agnelli G, Corea F, Ageno W, Alberti A, Lanari A, Caso V, Micheli S, Bertolani L, Venti M, Palmerini F, Biagini S, Comi G, Previdi P, Silvestrelli G. Early hemorrhagic transformation of brain infarction: rate, predictive factors, and influence on clinical outcome results of a prospective multicenter study. *Stroke* 2008;**39**:2249–2256.
63. Prodan C, Stoner J, Cowan L, Dale G. Lower coated-platelet levels are associated with early hemorrhagic transformation in patients with non-lacunar brain infarction. *J Thromb Haemost* 2010;**8**:1185–1190.
64. Kerr EA, Heister M, Krein SL, Kabeto M, Langa KM, Weir D, Piette JD. Beyond comorbidity counts: how do comorbidity type and severity influence diabetes patients' treatment priorities and self-management? *J Gen Intern Med* 2007;**22**:1635–1640.
65. Turner BJ, Hollenbeck CS, Weiner M, Ten Have T, Tang SS. Effect of unrelated comorbid conditions on hypertension management. *Ann Intern Med* 2008;**148**:578–586.
66. Silverberg DS, Wexler D, Iaina A, Schwartz D. The interaction between heart failure and other heart diseases, renal failure, and anemia. *Semin Nephrol* 2006;**26**:296–306.
67. Steg PG, James SK, Atar D, Badano LP, Lundqvist CB, Borger MA, Di Mario C, Dickstein K, Ducrocq G, Fernandez-Aviles F, Gershlick AH, Giannuzzi P, Halvorsen S, Huber K, Juni P, Kastrati A, Knuuti J, Lenzen MJ, Mahaffey KW, Valgimigli M, van 't Hof A, Widimsky P, Zaher D. ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J* 2012;**33**:2569–2619.
68. Amsterdam EA, Wenger NK, Brindis RG, Casey DE, Ganiats TG, Holmes DR, Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ. 2014 AHA/ACC guideline for the management of patients with non–ST-elevation acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014;**64**:e139–e228.