

**Muscle Glycogen Utilisation during an Australian Rules Football Game**

**Submission Type:** Case-Study

Harry E Routledge<sup>1,2</sup>, Jill J Leckey<sup>3</sup>, Matt J Lee<sup>4</sup>, Andrew Garnham<sup>3</sup>,  
Stuart Graham<sup>2</sup>, Darren Burgess<sup>2</sup>, Louise M Burke<sup>3</sup>, Robert M  
Erskine<sup>1,5</sup>, Graeme L Close<sup>1</sup> and James P Morton<sup>1</sup>

<sup>1</sup>Reserach Institute for Sport and Exercise Sciences  
Liverpool John Moores University  
Byrom Street Campus  
Liverpool  
L3 3AF  
UK

<sup>2</sup>Port Adelaide Football Club  
Allan Scott Headquarters  
Alberton Oval  
Brougham Place  
Alberton  
SA 5014  
Australia

<sup>3</sup>Exercise and Nutrition Research Program  
Mary MacKillop Institute for Health Research  
Australian Catholic University  
Melbourne  
Victoria, Australia

<sup>4</sup> College of Sport and Exercise Science  
Victoria University  
Melbourne  
Victoria  
Australia

<sup>5</sup>Institute of Sport, Exercise and Health,  
University College London,  
London,  
United Kingdom

**Address for correspondence:**

Dr James Morton  
Research Institute for Sport and Exercise Sciences  
Liverpool John Moores University  
Tom Reilly Building  
Byrom St Campus  
Liverpool  
L3 3AF  
United Kindgdom  
Email: [J.P.Morton@ljmu.ac.uk](mailto:J.P.Morton@ljmu.ac.uk)  
Tel: +44 151 904 6233

56  
57 **Running Head:** Glycogen use in AF match play  
58 **Abstract Word Count:** 248  
59 **Text Only Word Count:** 1492  
60 **Number of Figures and Tables:** 1 Table  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105

## Abstract

**Purpose:** To better understand the carbohydrate (CHO) requirement of Australian Football (AF) match play by quantifying muscle glycogen utilisation during an in-season AF match.

**Methods:** After a 24 h CHO loading protocol of 8 g/kg and 2 g/kg in the pre-match meal, two elite male forward players had biopsies sampled from *m. vastus lateralis* before and after participation in a South Australian Football League game. Player A (87.2kg) consumed water only during match play whereas player B (87.6kg) consumed 88 g CHO via CHO gels. External load was quantified using global positioning system technology.

**Results:** Player A completed more minutes on the ground (115 vs. 98 min) and covered greater total distance (12.2 vs. 11.2 km) than Player B, though with similar high-speed running (837 vs. 1070 m) and sprinting (135 vs. 138 m), respectively. Muscle glycogen decreased by 66% in Player A (Pre-: 656, Post-: 223 mmol·kg<sup>-1</sup> dw) and 24% in Player B (Pre-: 544, Post-: 416 mmol·kg<sup>-1</sup> dw), respectively.

**Conclusion:** Pre-match CHO loading elevated muscle glycogen concentrations (i.e. >500 mmol·kg<sup>-1</sup> dw), the magnitude of which appears sufficient to meet the metabolic demands of elite AF match play. The glycogen cost of AF match play may be greater than soccer and rugby and CHO feeding may also spare muscle glycogen use. Further studies using larger sample sizes are now required to quantify the inter-individual variability of glycogen cost of match play (including muscle and fibre-type specific responses) as well examine potential metabolic and ergogenic effects of CHO feeding.

**Keywords:** carbohydrate loading, vastus lateralis, AF, high speed running

## Introduction

Invasive team sports such as soccer<sup>1,2</sup>, rugby league<sup>2</sup> and Australian Football (AF)<sup>2,3</sup> are characterised by high-intensity (>19.8km.h<sup>-1</sup>) intermittent activity profiles. Given the duration of activity (i.e. 80-120 minutes) and high-intensity intermittent profiles, muscle glycogen is considered as the predominant energy substrate to support the metabolic demands of match play<sup>4,5</sup>. In relation to soccer, it has been reported that match-play in elite Danish soccer players<sup>5</sup> depletes muscle glycogen concentration by a magnitude of 50% and requires a glycogen cost of approximately 200 mmol.kg<sup>-1</sup> dw. We also observed similar absolute glycogen utilisation and relative depletion rates in English professional rugby league players during competitive match play<sup>6</sup>. Nonetheless, in the absence of a controlled carbohydrate (CHO) loading protocol, it is noteworthy that pre-match muscle glycogen concentrations in team sport athletes may range from 300-600 mmol.kg<sup>-1</sup> dw<sup>7</sup> and that approximately 50% of muscle fibres are classified as empty or partially empty after soccer match play, thus having potential implications for high-intensity physical performance late in the match. Indeed, in relation to physical performance, it was recognised as early as the 1970s<sup>8</sup> that commencing match play with reduced pre-exercise muscle concentration (i.e. < 200 mmol.kg<sup>-1</sup> dw) reduces total distance covered by 25% when compared with higher pre-match muscle glycogen availability (i.e. > 400 mmol.kg<sup>-1</sup> dw).

Given the longer duration and greater proportion of time spent at high-intensity workloads in AF match play<sup>2,3</sup> versus both soccer<sup>1,2</sup> and rugby<sup>2</sup>, it could be suggested that the CHO requirements for AF players are increased accordingly. Indeed, high velocity running was reported to be significantly greater in AF (1322m) versus rugby league (327m) and soccer (517m)<sup>2</sup>. To the authors' knowledge, however, no researchers have yet quantified the muscle glycogen cost of AF match play in elite players. The aim of the present case-study was to quantify glycogen use in *m. vastus lateralis* from two elite male AF players after participation in a South Australian Football League game.

## Methods

### Subjects

Two male forward players from a South Australian Football League (SANFL) list (see Table 1) volunteered to take part in the study. The study was conducted according to the

206 Declaration of Helsinki and was approved by the Human  
207 Research Ethics Committee of Australian Catholic University  
208 Melbourne.

209  
210

## 211 **Design**

212 In a case-study design, muscle biopsies were obtained from *m.*  
213 *vastus lateralis* before and after participation in a competitive  
214 SANFL league game undertaken in August 2017.  
215 Quantification of external loading during match play was  
216 assessed via global positioning system (GPS) technology.

217

218 **Dietary Controls.** In the day prior to match play, each player  
219 consumed a prescribed CHO loading diet providing CHO,  
220 protein and fat intakes corresponding to 8 g/kg, 2 g/kg and 1  
221 g/kg body mass, respectively, based on contemporary  
222 guidelines of Thomas and colleagues<sup>9</sup>. At 4 hours prior to  
223 match play, each player also consumed a pre-match meal  
224 providing 2 g/kg CHO, 0.2 g/kg protein and 0.3 g/kg fat.  
225 During match play, players were able to consume fluids and  
226 sports foods at opportunities that were both predictable  
227 (between quarters) and random (rotation on the bench, access to  
228 trainer on the ground when the ball is not in play). On such  
229 occasions; Player A consumed water only whereas Player B  
230 also consumed an additional 88 g of CHO in the form of four  
231 isotonic CHO gels (SiS GO Gels, Nelson, UK) consumed at the  
232 quarter time breaks (~20-minute intervals during the game).

233

234 **Muscle Biopsies.** Muscle biopsies (100 mg) were sampled  
235 from *m. vastus lateralis* at 60 minutes prior to the beginning of  
236 the game (i.e. prior to the warm-up period) and within 10  
237 minutes of completion of match play participation. Biopsies  
238 were obtained under local anesthesia (1% Xylocaine) using a  
239 Bergstrom biopsy needle and immediately frozen in liquid  
240 nitrogen for later analysis. Approximately 3 mg of freeze-dried  
241 muscle tissue was powdered and dissected free of all visible  
242 non-muscle tissue. Powdered muscle tissue was then mixed with  
243 250 µl of 2 M HCl, incubated at 95°C for 2 hours (agitated  
244 gently every 20 min), and then neutralized with 750µl of 0.66  
245 M NaOH. Glycogen concentration was subsequently assayed  
246 in triplicate via enzymatic analysis with flurometric detection.  
247 Muscle glycogen values were expressed as millimoles per  
248 kilograms dry weight (mmol·kg<sup>-1</sup> dw).

249

250 **GPS Analysis.** Both players wore a portable micro-technology  
251 device (Optimeye S5, Catapult Innovations, Melbourne,  
252 Australia) which recorded activity profile data. The portable  
253 device was worn inside a purpose built elastic vest, positioned  
254 across the upper back between the scapula. Each device was  
255 activated 30-minutes prior to the start of the game to allow

acquisition of satellite signal and lock (>8 satellites). Satellite data sampled at 10 Hz provided measures of duration, total distance, average speed and distance covered within four specific velocity bands corresponding to: jogging (2-3.9 m.s<sup>2</sup>), running (4-5.4 m.s<sup>2</sup>), high-speed running (5.5-6.9 m.s<sup>2</sup>) and sprinting (7 m.s<sup>2</sup>). These speed zones are similar to those previously reported in soccer<sup>1</sup> and allow for direct comparisons between football codes. At the conclusion of the match, data were downloaded and analyzed using (Openfield version 11.12, Catapult Innovations).

## Results

Subject characteristics, external workload, muscle glycogen utilisation and CHO intake are presented in Table 1. Individual data are presented for both Player A and B.

## Discussion

The aim of the present case-study was to quantify muscle glycogen utilisation during AF match play. We studied two elite male AF players during a competitive match from the South Australian Football League. Our data demonstrate that a standardised one day dietary CHO loading protocol of 8 g/kg and 2 g/kg in the pre-match meal elevates muscle glycogen concentration (i.e. >500 mmol.kg<sup>-1</sup> dw) to a magnitude that is sufficient to fuel the metabolic demands of AF match play.

In relation to the external loads reported here, we observed similar loading profiles to that previously reported in elite AF match play where larger sample sizes (e.g. 39 players) have been studied<sup>3</sup>. This is the case for parameters such as total distance, average speed, high-intensity running and sprinting. The between player differences in such parameters are also similar to those previously reported<sup>3</sup>. Despite only studying two forward players, we are confident that the external loads reported here are therefore representative of the customary loads experienced in AF match play from a wider sample of teams and players, though we acknowledge that load differences between playing positions are to be expected<sup>3</sup>.

In accordance with a higher workload, our data suggest that the glycogen cost of AF match play (e.g. Player A utilised 433 mmol.kg<sup>-1</sup> dw) may be greater than that reported in soccer<sup>5,7</sup> and rugby match play<sup>6</sup> where approximately 200 mmol.kg<sup>-1</sup> dw was utilised in both instances. Such differences in absolute glycogen cost are likely due to greater duration of activity and time spent in higher intensity threshold zones<sup>10</sup>. In contrast to Player A, Player B experienced less total glycogen use (138

mmol.kg<sup>-1</sup> dw). While such inter-individual variation in glycogen use may be due to differences in total distance covered, duration, pre-match muscle glycogen concentrations<sup>11</sup>, training status and muscle oxidative capacity<sup>12</sup>, it is noteworthy that Player B also consumed an additional 88 g of CHO during match play. As such, differences in glycogen use between players may also be due, in part, to a potential muscle glycogen sparing effect of CHO feeding, an effect that has been reported previously in *m. vastus lateralis* during running<sup>13</sup>. Indeed, the game characteristics of AF support a more aggressive approach to CHO intake during play than is reported by elite soccer players<sup>2</sup>, with opportunities for fuel replacement during scheduled breaks between quarters and time spent on the interchange bench. Nonetheless, we acknowledge that randomised control trials incorporating larger sample sizes are now required to verify any potential metabolic or ergogenic effect of CHO feeding during competitive match play. Additionally, glycogen use in specific muscles (e.g. gastrocnemius versus vastus lateralis), muscle fibre types and intra-cellular storage pools could also be quantified using transmission electron microscopy<sup>14,15</sup>. We also acknowledge that the glycogen utilisation observed here is also reflective of those activities undertaken during the warm-up period. As such, future studies could also sample biopsies in the minutes prior to match play and at the end of each quarter to further characterise both the total absolute use and rates of glycogen use as the match progresses.

## Practical Applications

Pre-match muscle glycogen concentration > 500 mmol.kg<sup>-1</sup> dw (as achieved via CHO loading of 8 g/kg) is sufficient to fuel the physical demands of elite forward AF match play. Given the apparently greater muscle glycogen cost of AF match-play compared to soccer and rugby, sport physiologists and nutritionists should ensure that AF players consume sufficient dietary CHO intake (likely > 6 g/kg) in the 24 hours prior to participation in match play.

## Conclusion

We provide novel data by reporting muscle glycogen utilisation in *m. vastus lateralis* from two elite male AF forward players during competitive match play. Our data suggest that total glycogen use is greater than that reported in elite players from other invasive team sports, such as soccer and rugby. Additionally, these data suggest that CHO loading with 8 g/kg body mass is sufficient to meet the metabolic demands of AF

356 match play and that previous suggestions of 10-12 g/kg body  
357 (for athletes involved in intermittent exercise > 90 minutes)  
358 mass may not be necessary for this population. Further studies  
359 are now required to quantify the inter-individual variability of  
360 glycogen use as well as examine any potential metabolic and  
361 ergogenic effects of CHO feedings during match play.

362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405



## References

1. Anderson L, Orme P, Di Michele R, Close GL, Morgans R, Drust B, Morton JP. Quantification of training load during one-, two- and three-game week schedules in professional soccer players from the English Premier League: implications for carbohydrate periodisation. *J Sports Sci.*2016;34(13):1250-1259.
2. Varley MC, Gabbett T, Aughey RJ. Activity profiles of professional soccer, rugby league and Australian football match play. *J Sports Sci.*2014;32(20):1858-1866.
3. Coutts AJ, Kempton T, Sullivan C, Bilsborough J, Cordy J, Rampinini E. Metabolic power and energetic costs of professional Australian Football match-play. *J Sci Med Sport.*2015;18(2):219-224.
4. Mujika I, Burke LM. Nutrition in team sports. *Ann Nutr Metab.*2010;57(2):26-35.
5. Gunnarsson TP, Bendiksen M, Bischoff R, Christensen PM, Lesvig B, Madsen K, Stephens F, Greenhaff P, Krstrup P, Bangsbo J. Effect of whey protein- and carbohydrate-enriched diet on glycogen resynthesis during the first 48 h after a soccer game. *Scand J Med Sci Sports.*2013;23(4):508-515.
6. Bradley WJ, Morehen JC, Haigh J, Clarke J, Donovan TF, Twist C, Cotton C, Shepard S, Cocks M, Sharma A, Impey SG, Cooper RG, MaClaren DP, Morton JP, Close GL. Muscle glycogen utilisation during Rugby match play: Effects of pre-game carbohydrate. *J Sci Med Sport.* 2016;19(12):1033-1038.
7. Krstrup, P, Mohr M, Steensberg A, Bencke, J, Kjaer M, Bangsbo J. Muscle and blood metabolites during a soccer game: implications for sprint performance. *Med Sci Sports Exerc.*2006;38(6):1165-1174.
8. Saltin B. Metabolic fundamentals in exercise. *Med Sci Sports.*1973;5(3):137-146.
9. Thomas DT, Erdman KA, Burke LM. Position of the academy of nutrition and dietetics, dietitians of canada,

and the american college of sports medicine: nutrition  
and athletic performance.*J Acad Nutr  
Diet.*2016;116(3):501-28.

10. Bangsbo J, Norregaard L, Thorsoe F. The effect of  
carbohydrate diet on intermittent exercise performance.  
*Int J Sports Med.*1992;13(2):152-157.
11. Wisbey B, Montgomery PG, Pyne DB, Rattray B.  
Quantifying movement demands of AFL football using  
GPS tracking.*J Sci Med Sport.*2010;13:531-536.
12. Hargreaves M, McConell G, Proietto J. Influence of  
muscle glycogen on glycogenolysis and glucose uptake  
during exercise in humans. *J Appl  
Physiol.*1995;78(1):288-292.
13. Holloszy JO, Coyle EF. Adaptations of skeletal muscle  
to endurance exercise and their metabolic  
consequences. *J Appl Physiol Respir Environ Exerc  
Physiol.*1984;56(4):831-838.
14. Tsintzas OK, Williams C, Boobis L, Greenhaff P.  
Carbohydrate ingestion and glycogen utilization in  
different muscle fibre types in man. *J Physiol.*1995;489  
:243-250.
15. Nielsen J, Krstrup P, Nybo L, Gunnarsson TP, Madsen  
K, Schroder HD, Bangsbo J, Ortenblad N. Skeletal  
muscle glycogen content and particle size of distinct  
subcellular localizations in the recovery period after a  
high-level soccer match. *Appl  
Physiol.*2012;112(10):3559-3367.

505

506

507 **Table 1** – Subject characteristics, external workload, muscle  
 508 glycogen utilisation and in-game CHO intake in Player A and B.

509

	<b>Player A</b>	<b>Player B</b>
<b>Playing Position</b>	Forward	Forward
<b>Age (years)</b>	22	27
<b>Body Mass (kg)</b>	87.2	87.6
<b>Height (m)</b>	1.88	1.82
<b>Warm Up Duration (min)</b>	12	12
<b>Distance (m)</b>	1478	1501
<b>High Speed Running (m) 5.5-6.9 m.s<sup>2</sup></b>	156	159
<b>Sprinting (m) &gt;7 m.s<sup>2</sup></b>	23	31
<b>Match Play Duration (min)</b>	115	98
<b>Number of Rotations</b>	2	4
<b>Average Speed (m/min)</b>	106	114
<b>Total Distance (m)</b>	12229	11182
<b>Walking (m) 0.1-1.9 m.s<sup>2</sup></b>	3801	2801
<b>Jogging (m) 2-3.9 m.s<sup>2</sup></b>	5231	4718
<b>Running (m) 4-5.4 m.s<sup>2</sup></b>	2180	2396
<b>High Speed Running (m) 5.5-6.9 m.s<sup>2</sup></b>	837	1070
<b>Sprinting (m) &gt;7 m.s<sup>2</sup></b>	135	138
<b>Pre-Match Glycogen (mmol.kg<sup>-1</sup> dw)</b>	656	544
<b>Post-Match Glycogen (mmol.kg<sup>-1</sup> dw)</b>	223	416
<b>Total Glycogen Utilisation (mmol.kg<sup>-1</sup> dw)</b>	433	138
<b>Total Exogenous CHO Consumed (g)</b>	0	88
<b>Exogenous CHO (g/h)</b>	0	54
<b>Exogenous CHO (g/min)</b>	0	0.9

510