

## PARV4 found in wild chimpanzee faeces - alternate route of transmission?

Kristýna Brožová<sup>a</sup>, David Modrý<sup>b,c,d</sup>, Eva Dadáková<sup>a</sup>, Mwanahamisi I. Mapua<sup>b</sup>, Alex K. Piel<sup>e,f</sup>, Fiona A. Stewart<sup>e,f</sup>, Vladimír Celer<sup>a</sup>, Kristýna Hrazdilová<sup>d,g</sup>

<sup>a</sup> Department of Infectious Diseases and Microbiology, University of Veterinary and Pharmaceutical Sciences Brno, Palackého tř. 1946/1, 612 42 Brno, Czech Republic

<sup>b</sup> Department of Pathological Morphology and Parasitology, University of Veterinary and Pharmaceutical Sciences Brno, Palackého tř. 1946/1, 612 42 Brno, Czech Republic

<sup>c</sup> Biology Centre, Institute of Parasitology, Czech Academy of Sciences, Branišovská 31, 370 05 České Budějovice, Czech Republic

<sup>d</sup> CEITEC-VFU, University of Veterinary and Pharmaceutical Sciences Brno, Palackého tř. 1946/1, 612 42 Brno, Czech Republic

<sup>e</sup> School of Natural Sciences and Psychology, Liverpool John Moores University, Byrom Street, Liverpool L33AF, UK

<sup>f</sup> Greater Mahale Ecosystem Research and Conservation Project (GMERC), Dar es Salaam, Tanzania

<sup>g</sup> Department of Virology, Veterinary Research Institute, Hudcova 296/70, 621 00 Brno, Czech Republic

**Corresponding author:** Kristýna Hrazdilová; [kristyna@hrazdilova.cz](mailto:kristyna@hrazdilova.cz), CEITEC-VFU, University of Veterinary and Pharmaceutical Sciences Brno, Palackého tř. 1946/1, 612 42 Brno, Czech Republic

### Abstract

Human parvovirus 4 (PARV4, family *Parvoviridae*, genus *Tetraparvovirus*) displays puzzling features, such as uncertain clinical importance/significance, unclear routes of transmission and discontinuous geographical distribution. The origin, or the general reservoir, of human PARV4 infection is unknown. We aimed to detect and characterize PARV4 virus in faecal samples collected from two wild chimpanzee populations and 19 species of captive non-human primates. We aimed to investigate these species as a potential reservoir and alternate route of transmission on the African continent. From almost 500 samples screened, a single

wild *Pan troglodytes schweinfurthii* sample tested positive. Full genome analysis, as well as single ORF phylogenies, confirmed species-specific PARV4 infection.

**Keywords:** PARV4, non-human primates, Africa, phylogeny

Human parvovirus 4 (PARV4, genus *Tetraparvovirus*, family *Parvoviridae*) displays puzzling features, such as uncertain clinical importance/significance, unclear routes of transmission and discontinuous geographical distribution. The first primate tetraparvovirus probably emerged in the 1980's together with the appearance of the HIV virus infection [1] and, so far, three phylogenetically distinct genotypes have been described in humans. Since first human PARV-4 description in 2005, the epidemiology data are growing and were recently reviewed by Matthews *et al.* [2]. Human PARV4 G1 and G2 are predominant in Europe, North America and Middle East [3–10], and are most often described in patients with a history of parenteral drugs application suffering from HIV, HCV or HBV infections [10, 11]. Genotype 2 is also reported from Asia [6, 12]. Human PARV4 G3 is distributed across Africa [13, 14] and exhibits unique features of transmissibility. It has been detected in diverse cohorts in the absence of other blood-borne viruses and/or intravenous drug/therapy application history [13–16], suggesting transmission routes other than parenteral infection. High frequencies of seropositivity in wild chimpanzee and gorilla serum samples also support the existence of another route of transmission of tetraparvoviruses in Africa [17].

The origin, or the general reservoir, of the human PARV4 is unknown. Three independent transmission events, perhaps from chimpanzees or other primate species, could be the source of the three genotypes [1, 17] in humans. A limited number of sequenced non-human primate tetraparvoviruses have been found to be species-specific for chimpanzees and colobus monkeys [17, 18]. Despite evidence of frequent exposure of African hunters to non-human primate bush meat, no direct evidence of cross-species parvovirus transmission has been found [18].

All previously described tetraparvovirus sequences from NHPs were obtained from either blood or bush meat samples [17, 18]. Our aim was to detect and characterize tetraparvoviruses in fecal samples collected from chimpanzees and other NHP with different levels of contact with humans, based on samples from (i) a wild, non-habituated eastern chimpanzee population in Issa Valley (Tanzania), known to be SIV positive [19], (ii) a wild, habituated eastern chimpanzee population from Kalinzu Forest Reserve (Uganda), and (iii) captive African non-human primates from Czech and Slovak zoos.

In total, 202 unidentified faecal samples were collected from eastern chimpanzees, with very limited contact with humans, in the Issa Valley (Figure 1) in Tanzania (between March 2012 and November 2013). The study site comprises approximately ~100 km<sup>2</sup> and is one of the driest and most open chimpanzee habitats, situated east of Lake Tanganyika, in western Tanzania [20]. The population density of Issa chimpanzees is estimated to be ~0.25 individuals/km<sup>2</sup> [21]. Further faecal samples (in total 123, 1-10 per individual) were collected from 42 identified individuals of habituated eastern chimpanzees (20 males, 22 females), with daily distant contact with researchers, in the Kalinzu Forest Reserve (Figure 1) [22] during April–July 2014. The forest reserve (~137 km<sup>2</sup>) is one of the three largest forest blocks in Uganda, being located on the eastern ridge of the western Rift Valley. The chimpanzee population density is estimated to be ~1.67 individuals/km<sup>2</sup> [23]. Captive primates, with daily intensive contact with keepers, were sampled in 13 Czech and Slovak zoological gardens; 25 *Pan troglodytes*, 10 *Gorilla gorilla* and 118 faecal samples from 17 different species other African primates were obtained [24] (see Supplementary material).

All faecal samples were preserved in RNAlater (Qiagen, Crawley, UK) and stored at -80°C until further use. DNA was isolated by SPRI (QIAquick PCR Purification Kit, Qiagen, Germany) according to manufacturer's instructions. DNA was screened by nested PCR based on a protocol published by Sharp *et al.* [17]. PCR products of the expected length of 295 nt were gel purified using QIAquick Gel Extraction Kit (Qiagen, Germany) according to manufacturer's instructions, cloned into pGEM®-T Easy Vector System (Promega, USA) and sequenced by Macrogen capillary sequencing services (MacrogenEurope, The Netherlands). Additional sets of primers located in conserved parts of primate tetraparvovirus and other tetraparvovirus genomes were designed to overlap each other. PCR products were purified, cloned and both strands sequenced as described above.

All obtained sequences were carefully edited using Geneious 11.0.2 [25] and compared with those available in GenBank by the BLASTn algorithm (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Sequence alignments were generated using the ClustalW algorithm. Phylogenetic trees were inferred by maximum likelihood method using IQ-TREE v. 1.6.beta4 [26]. A best-fit evolution model was then chosen based on the Bayesian information criterion (BIC) computed by ModelFinder [27]. Branch supports were assessed by the ultrafast bootstrap (UFBoot) approximation [28] and by SH-like approximate

likelihood ratio test (SH-aLRT) [29]. For detailed Material and Method see Supplementary material.

A total of 478 faecal samples was screened by nested PCR. Only a single sample from *P. t. schweinfurthii* (labelled U55) originating from the Issa Valley was found to be tetraparvovirus positive. The nearly whole U55 parvovirus genome of 4955 nt in length was sequenced (acc.no. MH215556). U55 chimpanzee PARV4 genome contains two main open reading frames (ORFs) and two additional small ORFs (ARF1 and 2) of unknown function, not observed in other members of the family *Parvoviridae* [14, 30]. The first main ORF of 1992 nt in length (U55 position of 59-2050) is located at the 5' end of the genome and encodes non-structural protein NS1. The second main ORF is located at the 3' end of the genome and consists of 2745 nt (positions 2147-4891), encoding two structural VP proteins (914 and 552 amino acids). Following current International Committee on Taxonomy of Viruses criteria, the U55 isolate belongs to the species *Primate tetraparvovirus 1* (with an amino acid sequence of NS1 protein identity 91.3-98.8% to human and chimpanzee genotypes within this species, Table).

In our phylogenetic analysis, the U55 sequence clusters together with other chimpanzee PARV4 sequences obtained from wild chimpanzees; *P. t. troglodytes* in Cameroon [17] and *P. t. verus* in Côte d'Ivoire [18] in both NS1 (Figure 2) and VP genes as well as in whole genome analysis (Supplementary material). This chimpanzee PARV4 clade is situated as a close outgroup to human genotypes with strong bootstrap support in all analyses (Figure 2, Supplementary material). Sequences published from colobus monkeys [18] formed separate clusters distinct from human as well as from chimpanzee PARV4 viruses. To further analyze the chimpanzee PARV4 genome, we performed a recombination analysis in Simplot software with reference strains of all human PARV4 genotypes and non-human primate PARV4 isolates other than chimpanzees. The analysis revealed no potential recombination sites that could have given rise to the novel chimpanzee PARV4 tetraparvovirus (data not shown).

Our study was inspired by growing evidence for alternate routes of transmission for PARV4 in sub-Saharan Africa [30–32]. Here we prove virus shedding is detectable in faeces through the detection of chimpanzee tetraparvovirus DNA in a single faecal sample collected from wild chimpanzees (*P. t. schweinfurthii*) from an SIV positive community in the Issa Valley (Tanzania). The low positivity rate observed among the Issa chimpanzee community can be explained by combined influence of low amount of excreted virus (detected only by nested, not simple PCR protocols) and shedding of the virus in limited time. Human PARV4 has been reported to be detected from faeces in a few cases, however, its shedding is very probably

intermittent or limited in time [13]. Detection of tetraparvoviruses in fecal samples and swabs was previously described in humans and NHP exclusively in sub-Saharan Africa [17, 18], suggesting this is either a unique characteristic of virus genotypes limited to Africa, or the impact of other factors unique to Africa, e.g. co-infection with other viruses/parasites (e.g. GI helminths), affecting tetraparvovirus shedding into the intestinal lumen. Repeated finding of PARV-4 virus in faeces is the first prerequisite to the faecal-oral route of transmission which deserves more attention as it implies a significant role in PARV4 infection spread in a period of acute infection in communities sharing territory and habits.

Phylogenetic and p-distance analyses of chimpanzee tetraparvovirus U55 described here place it along with sequences previously derived from *P. t. troglodytes* [17] and *P. t. verus* [18].

This fact, together with the lack of recombination sites detected in the genome proves the co-evolution of chimpanzee PARV-4 with their hosts, supporting opinion about host specificity of tetraparvoviruses [18]. Importantly, existence of chimpanzee-specific clade of PARV-4 suggests limited or nil possibility of zoonotic transfer of PARV-4 from chimpanzees to humans.

Complementary sets of blood/ faecal/nasal/other samples both from humans and non-human primates would be necessary to definitively address whether alternate routes of transmission of African tetraparvovirus occur and whether there is an influence of geographically restricted factors on the epidemiology of resulting infections.

## Acknowledgement

We thank the Tanzanian Wildlife Research Institute (TAWIRI) and Tanzanian Commission for Science and Technology (COSTECH) for permission to conduct research in Tanzania. This research was carried out under the project CEITEC 2020 (LQ1601) with financial support from the Ministry of Education, Youth and Sports of the Czech Republic under the National Program of Sustainability II, by project LO1218 with financial support from the Ministry of Education, Youth and Sports of the Czech Republic under the NPU I program, and further co-financed from the European Social Fund and the state budget of the Czech Republic (project OPVK CZ.1.07/2.3.00/20.0300). We acknowledge a grant for the development of research organization (RVO: RO0516). Support for GMERC (formerly UPP) and ongoing work at Issa comes from the UCSD/Salk Center for Academic Research and Training in Anthropogeny (CARTA).

## Compliance with Ethical Standards

Funding: All grants funding this study are stated under Acknowledgment section.

Conflict of Interest: All authors declare that he/she has no conflict of interest.

Ethical approval: This article does not contain any studies with human participants or animals performed by any of the authors.

## References

1. Manning A, Willey SJ, Bell JE, Simmonds P (2007) Comparison of tissue distribution, persistence, and molecular epidemiology of parvovirus B19 and novel human parvoviruses PARV4 and human bocavirus. *J Infect Dis* 195:1345–1352 . doi: 10.1086/513280
2. Matthews PC, Sharp C, Simmonds P, Klennerman P (2017) Human parvovirus 4 ‘PARV4’ remains elusive despite a decade of study. *F1000Research* 6:82 . doi: 10.12688/f1000research.9828.1
3. Asiyabi S, Nejati A, Shoja Z, et al (2016) First report of human parvovirus 4 detection in Iran. *J Med Virol* 88:1314–1318 . doi: 10.1002/jmv.24485
4. Fryer JF, Delwart E, Hecht FM, et al (2007) Frequent detection of the parvoviruses, PARV4 and PARV5, in plasma from blood donors and symptomatic individuals. *Transfusion* 47:1054–1061 . doi: 10.1111/j.1537-2995.2007.01235.x
5. Matthews PC, Malik A, Simmons R, et al (2014) PARV4: an emerging tetraparvovirus. *PLoS Pathog* 10:e1004036 . doi: 10.1371/journal.ppat.1004036
6. Prakash S, Jain A, Seth A, et al (2015) Complete Genome Sequences of Two Isolates of Human Parvovirus 4 from Patients with Acute Encephalitis Syndrome. *Genome Announc* 3:e01472-14 . doi: 10.1128/genomeA.01472-14
7. Rosenfeldt V, Norja P, Lindberg E, et al (2015) Low prevalence of parvovirus 4 in HIV-infected children in Denmark. *Pediatr Infect Dis J* 34:761–762 . doi: 10.1097/INF.0000000000000642
8. Servant-Delmas A, Laperche S, Lionnet F, et al (2014) Human parvovirus 4 infection in low- and high-risk French individuals. *Transfusion* 54:744–745 . doi: 10.1111/trf.12512
9. Sharp CP, Lail A, Donfield S, et al (2010) NIH Public Access. *J Infect Dis* 200:1119–1125 . doi: 10.1086/605646.High

10. Yu X, Zhang J, Hong L, et al (2012) High prevalence of human parvovirus 4 infection in HBV and HCV infected individuals in Shanghai. PLoS One 7:e29474 . doi: 10.1371/journal.pone.0029474
11. Simmons R, Sharp C, McClure CP, et al (2012) Parvovirus 4 infection and clinical outcome in high-risk populations. J Infect Dis 205:1816–1820 . doi: 10.1093/infdis/jis291
12. Chen MY, Hung C-C, Lee K-L (2015) Detection of human parvovirus 4 viremia in the follow-up blood samples from seropositive individuals suggests the existence of persistent viral replication or reactivation of latent viral infection. Virol J 12:94 . doi: 10.1186/s12985-015-0326-0
13. Drexler JF, Reber U, Muth D, et al (2012) Human Parvovirus 4 in Nasal and Fecal Specimens from Children, Ghana. Emerg Infect Dis 18:1650–1653 . doi: 10.3201/eid1810.111373
14. Sharp CP, Gregory WF, Hattingh L, et al (2017) PARV4 prevalence, phylogeny, immunology and coinfection with HIV, HBV and HCV in a multicentre African cohort. Wellcome Open Res 2:26 . doi: 10.12688/wellcomeopenres.11135.1
15. Sharp CP, Vermeulen M, Nébié Y, et al (2010) Changing epidemiology of human parvovirus 4 infection in sub-Saharan Africa. Emerg Infect Dis 16:1605–1607 . doi: 10.3201/eid1610.101001
16. Lavoie M, Sharp CP, Pépin J, et al (2012) Human parvovirus 4 infection, Cameroon. Emerg Infect Dis 18:680–683 . doi: 10.3201/eid1804.110628
17. Sharp CP, LeBreton M, Kantola K, et al (2010) Widespread Infection with Homologues of Human Parvoviruses B19, PARV4, and Human Bocavirus of Chimpanzees and Gorillas in the Wild. J Virol 84:10289–10296 . doi: 10.1128/JVI.01304-10
18. Adlhoch C, Kaiser M, Loewa A, et al (2012) Diversity of Chimpanzees , and Prey Relationships. Emerg Infect Dis 18:859–862 . doi: http://dx.doi.org/10.3201/eid1805.111849
19. Rudicell RS, Piel AK, Stewart F, et al (2011) High Prevalence of Simian Immunodeficiency Virus Infection in a Community of Savanna Chimpanzees. J Virol 85:9918–9928 . doi: 10.1128/JVI.05475-11
20. Dadáková E, Brožová K, Piel AK, et al (2018) Adenovirus infection in savanna chimpanzees (Pan troglodytes schweinfurthii) in the Issa Valley, Tanzania. Arch Virol 163:191–196 . doi: 10.1007/s00705-017-3576-x
21. Hernandez-Aguilar RA (2009) Chimpanzee nest distribution and site reuse in a dry habitat:

implications for early hominin ranging. *J Hum Evol* 57:350–364 . doi:  
10.1016/j.jhevol.2009.03.007

22. Mapua MI, Petrželková KJ, Burgunder J, et al (2016) A comparative molecular survey of malaria prevalence among Eastern chimpanzee populations in Issa Valley (Tanzania) and Kalinzu (Uganda). *Malar J* 15:423 . doi: 10.1186/s12936-016-1476-2
23. Howard P. (1991) *Nature conservation in Uganda's Tropical Forest Reserves*. IUCN, Gland, Switzerland and Cambridge, UK
24. Hrazdilová K, Slaninková E, Brožová K, et al (2016) New species of Torque Teno miniviruses infecting gorillas and chimpanzees. *Virology* 487:207–214 . doi: 10.1016/j.virol.2015.10.016
25. Kearse M, Moir R, Wilson A, et al (2012) Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649 . doi: 10.1093/bioinformatics/bts199
26. Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32:268–274 . doi: 10.1093/molbev/msu300
27. Kalyaanamoorthy S, Minh BQ, Wong TKF, et al (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nat Methods* 14:587–589 . doi: 10.1038/nmeth.4285
28. Minh BQ, Nguyen MA, Von Haeseler A, et al (2013) Ultrafast approximation for phylogenetic bootstrap. *Mol Biol Evol* 30:1188–1195 . doi: 10.1093/molbev/mst024
29. Guindon S, Gascuel O (2003) A Simple, Fast, and Accurate Algorithm to Estimate Large Phylogenies by Maximum Likelihood. *Syst Biol* 52:696–704 . doi: 10.1080/10635150390235520
30. Simmonds P, Douglas J, Bestetti G, et al (2008) A third genotype of the human parvovirus PARV4 in sub-Saharan Africa. *J Gen Virol* 89:2299–302 . doi: 10.1099/vir.0.2008/001180-0
31. Longhi E, Bestetti G, Acquaviva V, et al (2007) Human parvovirus 4 in the bone marrow of Italian patients with AIDS. *AIDS* 21:1481–1483 . doi: 10.1097/QAD.0b013e3281e38558
32. Panning M, Kobbe R, Vollbach S, et al (2010) Novel human parvovirus 4 genotype 3 in infants, Ghana. *Emerg Infect Dis* 16:1143–1146 . doi: 10.3201/eid1607.100025

**Conflicts of interest:** none

**Table:** Distance pairwise comparison of nucleotide and amino acid (boldface) distances of NS1 gene among strains belonging to the species *Primate tetraparvovirus 1*; cpzPARV4 – strains originating from chimpanzee, colPARV4 – strain originating from *Colobus polykomos*

		PARV4 G1	PARV4 G2	PARV4 G3	cpzPARV4			colPARV4
		EU546204	EU546205	EU874248	JN798203	U55	HQ113143	JN798211
PARV4 G1	EU546204		8,99	7,99	18,02	17,82	18,62	35,59
PARV4 G2	EU546205	2,56		8,4	18,17	17,77	18	36,41
PARV4 G3	EU874248	2,85	2,7		17,83	18,08	18,06	35,35
cpzPARV4	JN798203	8,3	8,6	9,17		11,7	10,49	34,88
	U55	7,99	7,99	8,73	3,62		5,5	35,49
	HQ113143	8,43	8,43	9,17	1,34	3,15		35,36
colPARV4	JN798211	32,43	32,43	33,01	32,43	32,28	32,41	

**Figure 1:** Map of study sites: Issa Valley (Tanzania) and Kalinzu (Uganda)

**Figure 2:** Phylogenetic analysis of full-length coding sequences of NS1 of strains belonging to the species *Primate tetraparvovirus 1* by maximum likelihood method using IQTREE software. Herein described strain is highlighted in red. Branch supports are displayed as % from 1000 replicates from SH-aLRT/UFBoot tests. Three different hokoviruses (EU200677, EU200669, JF504699) used as an outgroup are not displayed. Scale bar indicates a number of nucleotide substitutions per site.

Figure




Figure2

