



LJMU Research Online

Hamilton, CJ, Mammarella, IC and Giofré, D

Autistic-like traits in children are associated with enhanced performance in a qualitative visual working memory task.

<http://researchonline.ljmu.ac.uk/id/eprint/9623/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Hamilton, CJ, Mammarella, IC and Giofré, D (2018) Autistic-like traits in children are associated with enhanced performance in a qualitative visual working memory task. *Autism Research*, 11 (11). pp. 1494-1499. ISSN 1939-3792

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

**Autistic-like traits in children are associated with enhanced performance in a
qualitative visual working memory task.**

Colin J. Hamilton^{*}, Irene C. Mammarella^{**}, and David Giofrè^{**}

^{*}Department of Psychology, Northumbria University, Newcastle upon Tyne, UK

^{**}Department of Developmental and Social Psychology, University of Padova, Padova, Italy

^{***}Department of Natural Sciences and Psychology, Liverpool John Moores University,
Liverpool, UK

Paper accepted: 28/08/2018

Acknowledgments. The authors are extremely grateful to Chris Neasham and Jasmine Hassan for the collection of this data.

Conflict of Interest. Colin Hamilton states that he has no conflict of interest in pursuing and publishing this research. Irene Mammarella states that she has no conflict of interest in pursuing and publishing this research. David Giofrè states that he has no conflict of interest in pursuing and publishing this research.

Abstract

Prior research has suggested that individuals with Autism Spectrum Disorders (ASD) demonstrate heterogeneity in cognitive efficacy, challenged executive resources but efficient visual processing. These contrasts lead to opposing predictions about visuospatial working memory competency in both ASD and the broader autism phenotype (BAP); compromised by constrained executive processes, but potentially scaffolded by effective visual representation. It is surprising therefore, that there is a paucity of visual working memory (VWM) research in both the ASD and BAP populations, which have focused upon the visual features of the to-be-remembered stimulus. We assessed whether individual differences in VWM were associated with autistic-like traits (ALTs) in the BAP. About 76 children carried out the Visual Just Noticeable Difference task, designed to measure high fidelity feature representation within VWM. ALTs were measured with the Children's Empathy Quotient and Systemizing Quotient. Analyses revealed a significant positive relationship between Systemizing and VWM performance. This complements ASD studies in visual processing and highlights the need for further research on the working memory—long-term memory interface in ASD and BAP populations.

Lay Summary: This study was interested in how well children with high levels of autistic-like traits (ALTs) carry out a task which involved memorizing, for brief time, the precise size of coloured shapes. The results suggested that children with high levels of ALTs performed the task relatively well. This finding is in contrast to many previous studies suggest that ALTs are associated with poor memory, and suggests that future research needs to look more finely at how individuals carry out these tasks.

Keywords: broader autism phenotype; children; cognition; visual working memory; long-term memory

Autistic-like traits in children are associated with enhanced performance in a qualitative visual working memory task.

Autism Spectrum Disorder (ASD) is characterized by sustained challenges in reciprocal communication, social interaction, and restricted behaviors or activities; these behaviors are present in early development and may limit everyday activities [DSM-5; American Psychiatric Association, 2013]. In recent years, there has been an increasing recognition that the characteristics associated with ASD form a continuum and are continuously distributed throughout a wider population (Constantino & Todd, 2003; Le Couteur et al., 1996). The Broader Autism Phenotype (BAP) refers to the subclinical possession of these characteristics. One key characteristic of this continuum conceptualization is that observations made within ASD can generate research questions within the BAP population, and vice versa (Wallace, Budgett, & Charlton, 2016).

Individuals with ASD frequently present with an uneven profile of cognitive efficacy (e.g., Shah & Frith, 1983). A major contrast lies in the cognitive processes of visual perception and executive function (Chouinard, Parkington, Clements, & Landry, 2015). Extensive research findings have emphasized efficient perceptual processing relating to “attention to detail” of visual stimuli (Baron-Cohen, 2009; Happé & Frith, 2006; Mottron, Dawson, Soulières, Hubert, & Burack, 2006). However, with executive and attentional control resources, the findings have indicated severe challenges. This is evidenced across a number of functions: verbal fluency/long-term memory (LTM) access difficulties (Demetriou et al., 2017), generic executive challenges (Geurts, de Vries, & van den Bergh, 2014), and is congruent with Frith’s (2012) suggestion of challenged top-down control processes. This also appears to be the case in the BAP population (Christ, Kanne, & Reiersen, 2010).

These contrasting competencies raise interesting questions about the efficacy of working memory in ASD and BAP populations. Working memory has been defined as the

limited amount of information held in memory when undertaking tasks such as comprehension, learning, and problem solving; a process containing domain-general executive attentional resources interacting with domain-specific maintenance processes (Cowan, 2014). Within this definition, working memory in these populations could experience discrete and opposing impacts: impaired by challenged executive attentional resources but scaffolded by domain-specific representations underpinned by efficient visual perceptual encoding. A first glance at the ASD working memory literature indicates that the challenged executive resources appear to have the greatest impact with Kercood, Grskovic, Banda, and Begeske (2014) and Wang et al. (2017) concluding that visuospatial working memory is compromised in the ASD population. However, both raise caveats in their conclusions, emphasizing that in the majority of studies, it was spatial working memory (SWM) which was being examined and was impaired.

Research has suggested that visual working memory (VWM) and SWM can be dissociated both at the cognitive level (Darling, Della Sala, & Logie, 2007; Della Sala, Gray, Baddeley, Allamano, & Wilson, 1999; Hamilton, Coates, & Heffernan, 2003; Logie, 1995, 2011; Logie & Pearson, 1997) and at the neural level (Bellgowan, Buffalo, Bodurka, & Martin, 2009; Konstantinou, Constantinidou, & Kanai, 2017). In addition, evidence has suggested that SWM is more demanding of executive resources (Logie, 2011; Rudkin, Pearson, & Logie, 2007; Vandierendonck, Kemps, Fastame, & Szmalec, 2004).

Consequently, one could argue that while SWM could be challenged in ASD, VWM on the other hand could be more efficient due to enriched visual perceptual representation. What is therefore surprising, given the emphasis upon efficient visual processes within ASD, is the relative paucity of VWM research within both the ASD and BAP populations. Cui, Gao, Chen, Zou, and Wang (2010) carried out a study with children with Asperger's syndrome and observed that in a task demanding memory for visual patterns, the Asperger group were

impaired in performance. However, Hamilton (2011, 2013) has argued that tasks such as the one employed in Cui et al., demand executive attentional control resources in the form of grouping, organization, and LTM underpinning (Unsworth, Fukuda, Awh, & Vogel, 2014). Consequently, performance in such VWM tasks could still be compromised due to executive resource impairment. Mammarella, Giofrè, Caviola, Cornoldi and Hamilton (2014) found in a small ASD sample that when the visual patterns were manipulated in order to afford opportunities for semantic underpinning from LTM, the ASD group was particularly challenged. This indicated that the source of the difficulty was not necessarily in the memory process per se, but in a compromised executive attentional scaffolding memory performance (Wang et al., 2017).

There is also a lack of VWM studies in the BAP population; one exception to this is the Richmond, Thorpe, Berryhill, Klugman, and Olson (2013) article. This study looked at the relationship between autistic-like traits (ALTs) as measured by the Autism Spectrum Questionnaire (Baron-Cohen, Wheelwright, Skinner, Martin, & Clubley, 2001) and VWM performance in adults. An important element of their protocol was the use of memory stimuli that were difficult to verbalize. This manipulation most likely precluded the requirement for attentional control in the form of recoding and recruitment of verbal LTM semantics. They found that participants high in ALT characteristics performed the task more effectively. Unfortunately, one limitation in their protocol was that a sequential presentation format was employed and prior research has indicated that the final (recency) item presented is likely to have a different underlying visual memory representation from the pre-recency items (Allen, Baddeley, & Hitch, 2014; Phillips & Christie, 1977). Consequently, it is uncertain which component(s) contributed to the high ALT advantage in the Richmond et al. study.

The present study recruited a child sample and employed a qualitative VWM paradigm which aimed to minimize the requirement for semantic underpinning of the

representation as it focuses upon the quality or fidelity of the memory stimulus being maintained. To the authors' knowledge, no prior research has investigated qualitative VWM performance in the BAP. A simultaneous presentation procedure derived from the Thompson et al. (2006) was employed. In this protocol, participants were exposed to a memory stimulus and, after a maintenance period of 1 s were shown a probe stimulus and had to determine whether this probe stimulus was larger or smaller compared to the initially encoded stimulus. The Children's Empathy Quotient and Systemizing Quotient (SQ) (Auyeung et al., 2009) were employed to identify systemizing characteristics in children. Baron-Cohen (2009) has argued that systemizing characteristics should be associated with attention to detail in perception and memory. Age and gender were recorded as control variables. It was predicted that children high in systemizing ALT characteristics would perform the qualitative VWM task more effectively.

Methods

Participants

The study involved 76 children (29 boys and 47 girls) recruited from within mainstream schools in the UK. Exclusion criteria were impaired visual acuity and the presence of any developmental disorder. The age, Systemizing Quotient (SQ) and visual memory scores are shown in Table 1.

Table 1. Mean (SD) Age, Systemizing Quotient Score, and Size JND Task Performance as a Function of Gender

	Participant characteristics		
	Age in months (<i>SD</i>)	SQ score (<i>SD</i>)	Size JND task correct (<i>SD</i>)
Girls	100.32	27.43	21.43
<i>n</i> = 47	(-15.09)	(-7.05)	(2.20)
Boys	102.24	31.21	21.41
<i>n</i> = 29	(-15.71)	(-8)	(2.40)
Cohen <i>d</i>	0.13	0.51	0.01
(95% CIs)	-0.34, 0.59	0.03, 0.97*	-0.47, 0.45

Note. CIs: confidence intervals for Cohen's *d*. The text in italics are statistical values. * $p < 0.05$

Materials

Children's Empathy Quotient and SQ Questionnaire (Auyeung et al., 2009). This ALT questionnaire was given to the children's parents/carers to complete; and is composed of 28 Systemizing items. Systemizing scores conventionally range from 0 to 56. With the current Systemizing data, *Cronbach's* $\alpha = 0.72$.

Size Just Noticeable Difference (JND). This task was adapted from Thompson et al. (2006) (Fig. 1A,B). The retinal misalignment of the memory and probe stimuli ensured that iconic memory did not contribute to task performance (Phillips, 1974). Red or blue elliptical and rectangular stimuli were employed. The maintenance duration was 1 s following conventional change detection protocols (Luck & Vogel, 1997, 2013). All size changes were in the horizontal dimension. In no trial was the shape change categorical, for example, elliptical to circular; across the 24 trials, changes in horizontal axis varied from 5% to 30%, in either a larger or smaller extent (Fig. 1B). The maximum score was 24. The internal reliability of the Size JND was $\alpha = 0.683$. All viewing was within the child's arm reach of the laptop keyboard. Prior research has indicated that the Size JND task significantly reduces the

demands upon executive attentional control (Hamilton, 2013; Phillips & Hamilton, 2001; Thompson et al., 2006). The Size JND task is categorized as a VWM task (Cornoldi & Vecchi, 2003; Mammarella, Borella, Pastore, & Pazzaglia, 2013; Mammarella, Pazzaglia, & Cornoldi, 2008) which requires the precise memory for the size of a shape. This is in contrast to tasks viewed as simultaneous-spatial such as the Visual Patterns Task (VPT, Della Sala et al., 1999), where participants have to remember stimulus elements presented across space in a simultaneous manner. Also, in contrast to spatial-sequential tasks where spatially discrete elements of stimulus to be remembered are presented in a temporal sequence, such as the Corsi Blocks task.




Figure 1. The size JND task protocol. A, The temporal sequence of the procedure; B, an example of the memory stimulus and examples of the alternative probe stimuli which could be employed with this stimulus.

General Procedure

Children carried out the tasks in a quiet location within their school, and in the presence of the research assistant. The task took 10–15 min to complete. Ethical consent was granted by the Department of Psychology, Northumbria University. Written informed consent

was obtained from the schools and the participants' families, along with oral assent from the children.

Statistical Analyses

Hierarchical regression was employed initially, with subsequent regression moderation analysis using PROCESS (Hayes, 2017).

Results

Hierarchical regression analysis was used to identify whether age, gender, and the SQ score significantly predicted participants' Size JND performance. The three predictors explained 19.7% of the variance in Size JND task performance, $F(3,72) = 5.878, p = 0.001$. Age significantly predicted Size JND performance ($\beta = 0.245, p = 0.033$), however gender did not ($\beta = -0.134, p = 0.229$). Having controlled for age and gender, the SQ score uniquely predicted Size JND performance ($\beta = 0.476, p < 0.001$), and in addition, uniquely accounted for 18.9% of the variance in the memory scores (Fig. 2). Additional PROCESS analysis revealed that neither gender ($p = 0.610$) nor age ($p = 0.433$) moderated the relationship between the Systemizing score and Size JND task scores




Figure 2. The relationship between Systemizing and Size JND visual working memory task performance.

Discussion

This study investigated VWM performance and its association with Systemizing characteristics in children. A qualitative task was employed where participants had to briefly remember the precise size of the memory stimulus; a task emphasizing the quality of the memory representation, rather than the quantity of information held in working memory. Given the executive challenges found in ASD and BAP population it was anticipated that Size JND task demands would minimize recruitment from executive attentional control resources and thus reveal an advantage in the task for high Systemisers. This was indeed the case, with the SQ accounting for almost 19% of unique variance in visual memory performance. The present findings support the suggestion that observations of attention to detail in ASD (Baron-Cohen, 2009; Frith, 2012; Mottron et al., 2006) have positive implications for VWM task performance in the BAP population in a task context where the executive resource demand is attenuated.

The present study with children reveals a similar pattern of results to that of Richmond et al. (2013) study with adults in that higher ALTs were associated with improved VWM task performance. However, it should be noted that this earlier study by Richmond et al. employed the AQ measure to assess attention to detail and imagination autistic-like characteristics (Baron-Cohen et al., 2001). The VWM stimuli were random object shapes designed to preclude the recruitment of verbal semantics. Thus, the Richmond et al. study looked at object representation as opposed to the requirement of the retention of a surface visual feature (shape) in this current study (Dent, 2012). It is possible that the Richmond et al. and the current study found an advantage for individuals high in autistic-like characteristics because the two task protocols precluded the use of long-term memory verbal semantics in order to underpin task performance. Whereas in VWM tasks known to provide opportunities

to recruit LTM verbal semantics, tasks such as the VPT, ASD individuals seem unable to access and/or recruit these verbal semantics (Mammarella et al., 2014).

The present finding with its contrast to earlier SWM and executive resource findings has major implications for learning in both the ASD and BAP populations. The contrast between tasks demanding high quality VWM representation versus tasks demanding executive attentional control; tasks requiring “effort after meaning” (Frith, 2012). Given the key importance of the working memory—executive attention—long-term memory inter-face for education and learning (Cowan, 2014; Swanson & McMurren, 2018); future VWM research should be directed at understanding where, within this interface, individuals with ASD and/or high levels of systemizing characteristics have challenges in this complex interaction process. Clarification of this process in ASD may lead to the structuring of more effective learning environments, and putatively more effectively designed working memory intervention (de Vries, Prins, Schmand, & Geurts, 2015).

In conclusion, this study investigated the relationship between systemizing characteristics in children within the BAP population and performance in the qualitative Size JND VWM task. The results suggest a strong positive relationship between systemizing characteristics and the Size JND performance. This relative competence associated with ALTs is consistent with other VWM research in adults but in contrast to ASD VWM research indicating a compromised access to, and/or retrieval of, long-term memory verbal semantics. There is a need for future ASD and BAP working memory research to articulate in more detail the interface between working memory and long-term memory resources, and identify the relevant attentional control processes which are crucial for learning and education.

References

- Allen, R. J., Baddeley, A. D., & Hitch, G. J. (2014). Evidence for two attentional components in visual working memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *40*(6), 1499–1509. doi:10.1037/xlm0000002.
- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). Washington, DC: American Psychiatric Pub Incorporated.
- Auyeung, B., Wheelwright, S., Allison, C., Atkinson, M., Samarawickrema, N., & Baron-Cohen, S. (2009). The children's empathy quotient and systemizing quotient: Sex differences in typical development and in autism spectrum conditions. *Journal of Autism and Developmental Disorders*, *39*(11), 1509–1521. doi:10.1007/s10803-009-0772-x.
- Baron-Cohen, S. (2009). Autism: The empathizing-systemizing (E-S) theory. *Annals of the New York Academy of Sciences*, *1156*, 68–80. doi:10.1111/j.1749-6632.2009.04467.x.
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The autism-spectrum quotient (AQ): Evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*, *31*(6), 603–603.
- Bellgowan, P. S. F., Buffalo, E. A., Bodurka, J., & Martin, A. (2009). Lateralized spatial and object memory encoding in entorhinal and perirhinal cortices. *Learning & Memory*, *16*, 433–438.
- Chouinard, P., Parkington, K., Clements, B., & Landry, O. (2015). Dissociable visual perception and executive functioning processes in typically developing adults with varying degrees of autistic-like characteristics. *Frontiers Human Neuroscience*, *9*. doi:10.3389/conf.fnhum.2015.217.00109.

- Christ, S. E., Kanne, S. M., & Reiersen, A. M. (2010). Executive function in individuals with subthreshold autism traits. *Neuropsychology*, *24*(5), 590–598. doi:10.1037/a0019176.
- Constantino, J. N., & Todd, R. D. (2003). Autistic traits in the general population - A twin study. *Archives of General Psychiatry*, *60*(5), 524–530. doi:10.1001/archpsyc.60.5.524.
- Cornoldi, C., & Vecchi, T. (2003). Visuospatial working memory and individual differences. Hove, England: Psychology Press.
- Cowan, N. (2014). Working memory underpins cognitive development, learning, and education. *Educational Psychology Review*, *26*(2), 197–223. doi:10.1007/s10648-013-9246-y.
- Cui, J. F., Gao, D. G., Chen, Y. H., Zou, X. B., & Wang, Y. (2010). Working memory in early-school-age children with Asperger's syndrome. *Journal of Autism and Developmental Disorders*, *40*(8), 958–967.
- Darling, S., Della Sala, S., & Logie, R. H. (2007). Behavioural evidence for separating components within visuo-spatial working memory. *Cognitive Processing*, *8*(3), 175–181.
- de Vries, M., Prins, P. J. M., Schmand, B. A., & Geurts, H. M. (2015). Working memory and cognitive flexibility-training for children with an autism spectrum disorder: A randomized controlled trial. *Journal of Child Psychology and Psychiatry*, *56*(5), 566–576.
- Della Sala, S., Gray, C., Baddeley, A., Allamano, N., & Wilson, L. (1999). Pattern span: A tool for unwelding visuo-spatial memory. *Neuropsychologia*, *37*(10), 1189–1199.
- Demetriou, E. A., Lampit, A., Quintana, D. S., Naismith, S. L., Song, Y. J. C., Pye, J. E., ... Guastella, A. J. (2017). Autism spectrum disorders: A meta-analysis of executive function. *Molecular Psychiatry*, *23*, 1198–1204. doi:10.

1038/mp.2017.75.

- Dent, K. (2010). Dynamic visual noise affects visual short-term memory for surface color, but not spatial location. *Experimental Psychology*, *57*(1), 17–26.
- Frith, U. (2012). Why we need cognitive explanations of autism. *The Quarterly Journal of Experimental Psychology*, *65*(11), 2073–2092.
- Geurts, H. M., de Vries, M., & van den Bergh, S. F. W. M. (2014). *Executive functioning theory and autism*. In Handbook of executive function. New York: Springer.
- Hamilton, C., Coates, R., & Heffernan, T. (2003). What develops in visuo-spatial working memory development? *European Journal of Cognitive Psychology*, *15*(1), 43–69.
- Hamilton, C. J. (2011). The nature of visuospatial representation within working memory. In A. Vandierendonck & A. Smalec (Eds.), *Spatial working memory*. Hove: Psychology Press.
- Hamilton, C. J. (2013). The development of visuo-spatial working memory in children. In H. S. Clair-Thompson (Ed.), *Working memory: Developmental differences, component processes and improvement mechanisms*. New York: NOVA.
- Happé, F., & Frith, U. (2006). The weak coherence account: Detail-focused cognitive style in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, *36*(1), 5–25.
- Hayes, A. F. (2017). *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach (2nd ed.)*. New York: Guildford Press.
- Kercood, S., Grskovic, J. A., Banda, D., & Begeske, J. (2014). Working memory and autism: A review of literature. *Research in Autism Spectrum Disorders*, *8*(10), 1316–1332.
- Konstantinou, N., Constantinidou, F., & Kanai, R. (2017). Discrete capacity limits and neuroanatomical correlates of visual short-term memory for objects and spatial locations. *Human Brain Mapping*, *38*(2), 767–778. doi:10.1002/ hbm.23416.

- Le Couteur, A., Bailey, A., Goode, S., Pickles, A., Robertson, S., Gottesman, I., & Rutter, M. (1996). A broader phenotype of autism: The clinical spectrum in twins. *Journal of Child Psychology and Psychiatry*, 37(7), 785–801.
- Logie, R. H. (1995). Visuo-spatial working memory. Hove: Lawrence Erlbaum Associates Ltd.
- Logie, R. H. (2011). The functional organization and capacity limits of working memory. *Current Directions in Psychological Science*, 20(4), 240–245.
doi:10.1177/0963721411415340.
- Logie, R. H., & Pearson, D. G. (1997). The inner eye and the inner scribe of visuo-spatial working memory: Evidence from developmental fractionation. *European Journal of Cognitive Psychology*, 9(3), 241–257.
- Luck, S. J., & Vogel, E. K. (1997). The capacity of visual working memory for features and conjunctions. *Nature*, 390(6657), 279–281.
- Luck, S. J., & Vogel, E. K. (2013). Visual working memory capacity: From psychophysics and neurobiology to individual differences. *Trends in Cognitive Sciences*, 17(8), 391–400.
- Mammarella, I. C., Borella, E., Pastore, M., & Pazzaglia, F. (2013). The structure of visuospatial memory in adulthood. *Learning and Individual Differences*, 25, 99–110.
- Mammarella, I. C., Giofrè, D., Caviola, S., Cornoldi, C., & Hamilton, C. (2014). Visuospatial working memory in children with autism: The effect of a semantic global organization. *Research in Developmental Disabilities*, 35(6), 1349–1356.
- Mammarella, I. C., Pazzaglia, F., & Cornoldi, C. (2008). Evidence for different components in children's visuospatial working memory. *British Journal of Developmental Psychology*, 26(3), 337–355.
- Mottron, L., Dawson, M., Soulières, I., Hubert, B., & Burack, J. (2006). Enhanced

perceptual functioning in autism: An update, and eight principles of autistic perception. *Journal of Autism and Developmental Disorders*, 36(1), 27–43.

Phillips, L. H., & Hamilton, C. (2001). The working memory model in adult aging research. In J. Andrade (Ed.), *Working memory in perspective* (pp. 101–125). Hove, UK: Psychology Press.

Phillips, W. A. (1974). Distinction between sensory storage and short-term visual memory. *Perception & Psychophysics*, 16(2), 283–290.

Phillips, W. A., & Christie, D. F. M. (1977). Components of visual memory. *Quarterly Journal of Experimental Psychology*, 29, 117–133.
doi:10.1080/00335557743000080.

Richmond, L. L., Thorpe, M., Berryhill, M. E., Klugman, J., & Olson, I. R. (2013). Individual differences in autistic trait load in the general population predict visual working memory performance. *The Quarterly Journal of Experimental Psychology*, 66(6), 1182–1195. doi:10.1080/17470218.2012.734831.

Rudkin, S. J., Pearson, D. G., & Logie, R. H. (2007). Executive processes in visual and spatial working memory tasks. *The Quarterly Journal of Experimental Psychology*, 60(1), 79–100.

Shah, A., & Frith, U. (1983). An islet of ability in autistic children: A research note. *Journal of Child Psychology and Psychiatry*, 24(4), 613–620.

Swanson, H. L., & McMurren, M. (2018). The impact of working memory training on near and far transfer measures: Is it all about fluid intelligence? *Child Neuropsychology*, 24(3), 370–395. doi:10.1080/09297049.2017.1280142.

Thompson, J. M., Hamilton, C. J., Gray, J. M., Quinn, J. G., Mackin, P., Young, A. H., & Ferrier, I. N. (2006). Executive and visuospatial sketchpad resources in euthymic bipolar disorder: Implications for visuospatial working memory

architecture. *Memory*, 14(4), 437–451. doi:10.1080/09658210500464293.

Unsworth, N., Fukuda, K., Awh, E., & Vogel, E. K. (2014). Working memory and fluid intelligence: Capacity, attention control, and secondary memory retrieval.

Cognitive Psychology, 71, 1–26.

Vandierendonck, A., Kemps, E., Fastame, M. C., & Szmalec, A. (2004). Working memory components of the Corsi blocks task. *British Journal of Psychology*, 95(1), 57–79.

Wallace, G. L., Budgett, J., & Charlton, R. A. (2016). Aging and autism spectrum disorder: Evidence from the broad autism phenotype. *Autism Research*, 9(12), 1294–1303.

Wang, Y., Zhang, Y. B., Liu, L. L., Cui, J. F., Wang, J., Shum, D. H. K., ... Chan, R. C. K. (2017). A meta-analysis of working memory impairments in autism spectrum disorders. *Neuropsychology Review*, 27(1), 46–61.