

Secondary distribution of injecting equipment obtained from needle and syringe programmes by people injecting image and performance enhancing drugs: England & Wales, 2012-15.

Rachel Glass ^a, Vivian D Hope ^{a,b}, Jacquelyn Njoroge ^a, Claire Edmundson^a, Josie Smith ^c, James McVeigh^b, John Parry ^a, Monica Desai ^a.

- a. National Infection Service, Public Health England, London, UK.
- b. Public Health Institute, Liverpool John Moores University, Liverpool, UK.
- c. Health Protection, Public Health Wales, Cardiff, UK.

Corresponding author:

Rachel Glass

Drug Use & Infections Team, HIV & STI Department

National Infection Service, Public Health England

61 Colindale Avenue, London, NW9 5EQ, UK.

Rachel.Glass@phe.gov.uk

Tel: 020 8327 7038

Keywords: Secondary distribution, Image and performance enhancing drugs, Needle and syringe programmes, Harm reduction, people who inject drugs

Abstract

Background

People who inject image & performance enhancing drugs (IPEDs) are often the largest group using needle and syringe programmes (NSPs) in the UK. NSP providers report these clients repeatedly collecting large amounts of equipment for others. The extent of secondary distribution of injecting equipment is unknown.

Methods

Data from national surveillance of people injecting IPEDs was used. Participants completed a questionnaire and provided a dried-blood spot sample. Data from two biennial surveys was combined; repeat participants were excluded. Self-reported data was used to explore the extent of secondary distribution.

Results

Of the participants 87% (467) reported NSP use: median age 31 years, 98% male. A third (34%, 157) reported collecting equipment for others. Of those collecting for others, 154 reported how many people they had collected for: 55% had collected for one person, 27% for 2-9 people, 5% for 10-19 and 13% for 20 or more (no difference by psychoactive drug use). Those vaccinated for hepatitis B were more likely (22% [15/68] vs 6% [5/86], $p=0.003$) and those reporting redness/swelling at an injection site were less likely to collect equipment for at least 20 others (8% [8/106] vs 25% [12/48], $p=0.003$). Overall, 154 people collected equipment for 639-1,569 people injecting IPEDs.

Conclusions

Secondary distribution of injecting equipment is common among those injecting IPEDs and using NSPs. Whilst not allowing for rotational collection within groups, our analysis suggests

that many of those injecting IPEDs are not in direct contact with NSPs. Innovation approaches for harm reduction interventions are needed.

Word count: 249/250

1. Introduction

Historically, the use of IPEDs, such as anabolic steroids, had predominately been for athletic purposes (Begley et al., 2017). In recent years the range of drugs and prevalence of use for both aesthetic and athletic purposes has increased in England and Wales (Bates et al., 2014; Begley et al., 2017; Home Office, 2016). UK needle and syringe programmes (NSP) often report seeing more people who inject IPEDs than people who inject psychoactive drugs (Bates et al., 2014; McVeigh and Begley, 2017). Increased NSP use by those injecting IPEDs has been reported in other high income countries, with international debate concerning adequate service provision (Iversen et al., 2016).

Appropriate harm reduction interventions are vital to prevent adverse health outcomes, including blood borne viral (BBV) and skin and soft tissue infections (SSTI); 1 in 20 people who inject IPEDs in the UK are hepatitis C antibody positive and 1 in 7 report SSTI symptoms (Hope et al., 2013; Public Health England, July 2015). UK policy is to provide free needles and syringes to all people who inject drugs (PWID) and to support the return of used equipment (NICE, 2014b). The IPED injecting population is receptive to NSPs (NICE, 2014b). However, due to the cyclical nature of IPED use (Chandler and J.McVeigh, 2013), involving planned periods of use and non-use, NSP attendance among people using IPEDs can be less frequent and more seasonal compared to those injecting psychoactive drugs, with IPED injectors often collecting large quantities of equipment (McVeigh et al., 2003).

Qualitative research indicates secondary distribution is occurring, where people who inject IPEDs collect injecting equipment for distribution through their peer networks (McVeigh et al., 2007). UK guidance recommends that services should not discourage secondary distribution, but should encourage its recipients to personally attend NSPs for harm reduction interventions (NICE, 2014b). Among people who inject psychoactive drugs, secondary distribution is evident (Craine et al., 2010; De et al., 2008; Lorvick et al., 2006), with those reporting this as their primary source of clean equipment being younger and less

experienced, and so probably more vulnerable, PWID (Craine et al., 2010). The extent of peer-distribution within IPED networks is unclear, but could be considerable with the recipients potentially being a vulnerable population not reached by current interventions.

We estimate the current extent and nature of the secondary distribution of injecting equipment from English and Welsh NSP by people injecting IPEDs.

2. Methodology

Data from a national cross-sectional survey was used. As part of the Unlinked Anonymous Monitoring (UAM) Survey of PWID a biennial sub-survey of people injecting IPEDs was undertaken. Methodological details have been previously reported (Hope et al., 2013). In summary, people who have ever injected IPEDs were recruited through sentinel collaborating NSPs or outreach services. Participants completed a questionnaire and provided a dried blood spot sample, which was tested for antibodies against HIV, hepatitis B and hepatitis C (Hope et al., 2016; Hope et al., 2017). The survey has multi-site ethical approval from NHS Health Research Authority (MREC/98/2/51).

Data from two survey waves, covering the period 2012-2015 from 29 sentinel services were used. Participants in the second wave who reported participating in the first wave were excluded. Exploratory analysis found missing data in binary variables did not differ in the outcome variable to the non-affirmative responses, so missing data were recoded as such.

Descriptive and comparative analyses were conducted, using Pearson's Chi-squared test, to examine bivariate associations ($p < 0.05$) between the outcome variable (collecting equipment for others) with covariates (demographics, injecting practices, intervention uptake and BBV status).

Participants reporting NSP use were asked whether they had collected injecting equipment for others, and how many additional people they routinely collected for, categorised as: 1, 2-9, 10-19 and 20+ others. To assess the extent of onward distribution sensitivity analysis was

conducted, producing a minimum, mid-point and maximum estimate. Bivariate associations were examined for those reporting extensive onward distribution (to ≥ 20 others) and those who have ever injected IPEDs and psychoactive substances, with covariates previously described. Analyses were conducted in STATA 13.

3. Results

Of the 537 participants, 467 (87%) reported NSP use (Table 1). The demographic characteristics and injecting practises of those using NSP and those not were similar; reporting of redness or swelling at an injection site was, however, higher among the NSP users (Table 1). The median age of those reporting NSP use was 31 years, 98% were men, 1.5% (n=7) were living with HIV, 3.0% (n=14) and 5.1% (n=24) had ever had hepatitis B and hepatitis C respectively.

The IPEDs injected were those commonly reported, with 98% reporting use of anabolic steroids. One-third (34%, n=157) of those using NSPs reported usually collecting injecting equipment for others; the median age at first NSP use for those collecting for others was 26 years and their median time since first NSP use was five years (N=123).

Those collecting injecting equipment for others were more likely to report ever having redness, tenderness and swelling (68% vs 58%, $p=0.029$) and SSTI symptoms (21% vs 13%, $p=0.029$) at an injection site than those not collecting equipment for others (Table 1). Additionally, those reporting secondary distribution were more likely to also inject psychoactive substances (ever: 19% vs 11%, $p=0.021$; recently: 11% vs 4%, $p=0.011$) and report ever sharing a drugs vial (18% vs 9%, $p=0.002$) (Table 1). There was no difference in age.

3.1 Use of psychoactive drugs

Injecting psychoactive drugs in the past year was rare. Overall, 8% (n=36) of participants reported ever injecting heroin, 8% (n=36) cocaine, and 7% (n=32) speed. Participants who

had ever injected IPEDs and who also reported injecting psychoactive drugs during the past year (6%, $n=29$) were older, (median age 38 vs 30 years, $p=0.0013$) than those only injecting IPEDs ($n=438$), and more likely to have ever been incarcerated (69% [20/29] vs 22% [96/438], $p<0.001$), vaccinated for hepatitis B (83% [24/29] vs 38% [166/438], $p<0.001$), ever tested for hepatitis C (83% [24/29] vs 36% [157/438], $p<0.001$), reported symptoms of an SSTI (38% [11/29] vs 14% [63/438], $p=0.001$) and re-used injecting equipment (79% [23/29] vs 13% [55/438], $p<0.001$).

3.2 Extent of secondary distribution

Of the 154 who reported the number of other people they collected for, over half (55%) usually collected for one other person, around a quarter (27%) collected for two to nine other people, and 5% collected for ten to nineteen other people. The remaining 13% (4% of the overall sample) reported usually collecting for twenty or more other people (Table 2).

Overall, 154 people collected equipment for at least another 639 people injecting IPEDs. A less conservative approach, using a mid-point estimate, suggests collection for an additional 1,000 people injecting IPEDs, and at most an additional 1,569 people were collected for (Table 2).

Considering extensive onward distribution, those vaccinated for hepatitis B were more likely to collect equipment for at least 20 others (22% [15/68] vs 6% [5/86], $p=0.003$) and those ever reporting redness or swelling at an injection site were less likely to collect equipment for at least 20 others (8% [8/106] vs 25% [12/48], $p=0.003$). There were no significant differences with age, gender, BBV infection or the injection of psychoactive drugs.

Among those reporting secondary distribution there was no difference in the number of people they collected for, between those who had and those who had not injected psychoactive drugs in the preceding year ($p=0.456$).

4. Discussion

Our analysis confirms anecdotal reports that secondary distribution of injecting equipment obtained from NSPs by people who inject IPEDs is common, aligning with the single previous UK study (McVeigh et al., 2003). Moreover, our conservative estimate found more people received equipment indirectly through secondary distribution than directly from an NSP and, at most, the population indirectly supplied was around three times the size of our sample.

Although the extent of secondary distribution estimated has a wide range, it suggests that the group of people using IPEDs and not in direct contact with NSPs is probably large. Secondary distribution enables widespread needle delivery, allowing connection with hard to reach users (Kimergård and McVeigh, 2014) and is more effective in reducing BBV transmission than restrictive policies (Kerr et al., 2010). However, those obtaining injecting advice exclusively from peers could mimic their peer's behaviours. If so, our finding that collecting for others was associated with sharing equipment and SSTI symptoms is a concern. The HIV prevalence among those injecting IPEDs and accessing NSPs than in the general UK population is higher (Kirwan et al., 2016). Although prevalence among recipients of onward equipment distribution is unknown, those not in NSP contact may be at greater risk (Public Health England et al., 2016). Our findings, therefore, highlight the importance of alternative platforms and improved approaches for delivering harm reduction interventions to people injecting IPEDs.

One approach would be to utilise the secondary distribution process by engaging those collecting as peer educators (Bates et al., 2014; Marshall et al., 2015). This has previously been shown to be effective among people using psychoactive drugs (Anderson et al., 2003; Broadhead et al., 1998; Craine et al., 2006; Grund et al., 1992; Winyard, 2005), with distributors displaying a health-related altruism to protect others from harm (Bryant and Hopwood, 2009; Snead et al., 2003). Among people using IPEDs, knowledge around IPED

use is sourced primarily from peers, despite their knowledge sometimes being incomplete or inaccurate (Hanley Santos and Coomber, 2017).

Secondly, using social media and online networking for health promotion should be considered (Gold et al., 2012). Currently, people using IPEDs often rely on online forums and websites for information, which can be misleading or inaccurate and thus leading to harms (Bates et al., 2014). As IPEDs and injecting equipment are often sourced online (Andreas et al., 2014; Kimergard and McVeigh, 2014), NSPs could offer online ordering of injecting equipment, incorporating education to improve injecting practices, to engage with this group.

Thirdly, innovative outreach approaches within the IPED community could offer unique access, for delivery of both education and NSPs. The participants who reported extensive secondary distribution may have been collecting for specific gyms. They reported higher uptake of interventions and lower levels of poor injecting practices, indicative of an experienced sub-group with safer behaviours, thus potentially suitable candidates for delivering peer-led gym-based programmes. Such programmes do exist (NICE, 2014a) including informal arrangements with the local NSP (Kimergård and McVeigh, 2014), however obtaining owner consent can be a barrier (Bates et al., 2014).

Lastly, this study highlights the need for NSPs to appropriately address secondary distribution. Services, including pharmacy exchanges, should be willing and able to provide sufficient injecting equipment for additional onward distribution, alongside information in-line with current guidelines (NICE, 2014b).

People who inject IPEDs are mostly a discrete population from those who inject psychoactive drugs; however there is an overlap between these two groups. People injecting both IPEDs and psychoactive drugs were found to be older, report higher uptake of interventions, riskier injecting practices and more often collected equipment for others. A variety of factors may influence this: being older they may have been injecting for longer and

be more established in user networks, or they could be more comfortable attending NSPs due to greater exposure through psychoactive drug use, and therefore attend more often. However, the extent of secondary distribution was similar for people who had only injected IPEDs and those who had also injected psychoactive drugs. Compared to international studies among people who predominantly or exclusively inject psychoactive drugs, the proportion collecting for onward distribution in our study was lower (Bryant and Hopwood, 2009; Newland et al., 2016a; Tyndall et al., 2002), however psychoactive injectors often distribute in smaller networks or to partners (Bryant and Hopwood, 2009). Newland *et al.*, found a comparable proportion reporting large-scale distribution (16%) (Newland et al., 2016a) to our study (13%).

There are limitations to our study to consider. Firstly, we are unable to know whether the individuals collecting for multiple others are doing so on a rotational basis within their distribution network. Thus some of those being collected for may also use NSPs (Newland et al., 2016b). Additionally tertiary distribution, where clients may pass equipment onto an individual (potentially a training partner) who again distributes equipment to others, may be occurring and unaccounted for. We also have no information on those who only obtain needles from other sources, such as, through online purchases. Secondly, the data on NSP use relies on self-report and is therefore subject to recall bias. Despite the survey's anonymity, participants may be reluctant to disclose the extent of their redistribution due to concerns over a reduction in provision. Due to using discrete answer categories, our estimates of onward distribution include ambiguity. Finally, as information on the size and nature of this population is currently limited, we are unable to assess the representativeness of the sample and the generalizability of these findings.

5. Conclusion

Secondary distribution of injecting equipment obtained from NSPs is common among those injecting IPEDs. The number of people for whom equipment was collected was greater than the number participating in our study. Our study suggests that many, possibly a majority, of those injecting IPEDs are not in direct contact with NSPs. Those who were also injecting psychoactive drugs more often collected equipment for others. NSPs should target those collecting for others as potential peer educators for cascading harm reduction messages and explore alternative platforms for communicating safer injecting practices among people who inject IPEDs.

Words: 2050/2000

References

- Anderson, R., Clancy, L., Flynn, N., Kral, A., Bluthenthal, R., 2003. Delivering syringe exchange services through “satellite exchangers”: the Sacramento Area Needle Exchange, USA. *Int. J. Drug Policy*. 14, 461-463.
- Andreas, K., Jim, M., Simon, K., Torben, B., Allan, S., 2014. Online marketing of synthetic peptide hormones: poor manufacturing, user safety, and challenges to public health. *J. Drug Test. Anal.* 6, 396-398.
- Bates, G., Jones, L., McVeigh, J., 2014. Update of NICE Guidance PH18 Needle and Syringe Programmes (PIED Review). Centre for Public Health, Liverpool
- Begley, E., McVeigh, J., Hope, V., 2017. Image and Performance Enhancing Drugs: 2016 National Survey Results. Public Health Institute, Liverpool John Moores University.
- Broadhead, R.S., Heckathorn, D.D., Weakliem, D.L., Anthony, D.L., Madray, H., Mills, R.J., Hughes, J., 1998. Harnessing peer networks as an instrument for AIDS prevention: Results from a peer-driven intervention. *Public Health Rep.* 113, 42-57.
- Bryant, J., Hopwood, M., 2009. Secondary exchange of sterile injecting equipment in a high distribution environment: A mixed method analysis in south east Sydney, Australia. *Int. J. Drug Policy*. 20, 324-328.
- Chandler, M., J. McVeigh, 2013. Steroids and image enhancing drugs 2013 survey results. Centre for Public Health, John Moores University.
- Craine, N., Hickman, M., Parry, J.V., Smith, J., McDonald, T., Lyons, M., 2010. Characteristics of injecting drug users accessing different types of needle and syringe programme or using secondary distribution. *J. Public Health*. 32, 328-335.
- Craine, N., Mark Walker, A., Williamson, S., Bottomley, T., 2006. Reducing the risk of exposure to HCV amongst injecting drug users: Lessons from a peer intervention project in Northwest Wales. *Journal of Substance Use*. 11, 217-227.

- De, P., Cox, J., Boivin, J.F., Platt, R.W., Jolly, A.M., 2008. Social network-related risk factors for bloodborne virus infections among injection drug users receiving syringes through secondary exchange. *J. Urban Health*. 85, 77-89.
- Gold, J., Pedrana, A.E., Stoove, M.A., Chang, S., Howard, S., Asselin, J., Ilic, O., Batrouney, C., Hellard, M.E., 2012. Developing health promotion interventions on social networking sites: recommendations from The FaceSpace Project. *J. Med. Internet Res*. 14, e30.
- Grund, J.-P.C., Blanken, P., Adriaans, N.F.P., Kaplan, C.D., Barendregt, C., Meeuwssen, M., 1992. Reaching the Unreached: Targeting Hidden IDU Populations with Clean Needles via Known User Groups. *J. Psychoactive Drugs*. 24, 41-47.
- Hanley Santos, G., Coomber, R., 2017. The risk environment of anabolic–androgenic steroid users in the UK: Examining motivations, practices and accounts of use. *Int. J. Drug Policy*.
- Home Office, 2016. Drug misuse: Findings from the 2015/16 crime survey for England and Wales. London.
- Hope, V.D., Cullen, K.J., Smith, J., Jessop, L., Parry, J., Ncube, F., 2016. Is the recent emergence of mephedrone injecting in the United Kingdom associated with elevated risk behaviours and blood borne virus infection? *Euro Surveill*. 21.
- Hope, V.D., Iversen, J., Cullen, K.J., Parry, J.V., Maher, L., Ncube, F., 2017. Injection into the jugular vein among people who inject drugs in the United Kingdom: Prevalence, associated factors and harms. *Int. J. Drug Policy*. 46, 28-33.
- Hope, V.D., McVeigh, J., Marongiu, A., Evans-Brown, M., Smith, J., Kimergard, A., Croxford, S., Beynon, C.M., Parry, J.V., Bellis, M.A., Ncube, F., 2013. Prevalence of, and risk factors for, HIV, hepatitis B and C infections among men who inject image and performance enhancing drugs: a cross-sectional study. *BMJ Open*. 3, e003207.
- Iversen, J., Hope, V.D., McVeigh, J., 2016. Access to needle and syringe programs by people who inject image and performance enhancing drugs. *Int. J. Drug Policy*. 31, 199-200.

- Kerr, T., Small, W., Buchner, C., Zhang, R., Li, K., Montaner, J., Wood, E., 2010. Syringe Sharing and HIV Incidence Among Injection Drug Users and Increased Access to Sterile Syringes. *Am. J. Public Health.* 100, 1449-1453.
- Kimergard, A., McVeigh, J., 2014. Environments, risk and health harms: a qualitative investigation into the illicit use of anabolic steroids among people using harm reduction services in the UK. *BMJ Open.* 4, e005275.
- Kimergård, A., McVeigh, J., 2014. Variability and dilemmas in harm reduction for anabolic steroid users in the UK: a multi-area interview study. *Harm Reduction Journal.* 11, 19-19.
- Kirwan, P.D., Chau, C., Brown, A.E., Gill, O.N., Delpech, V.C., and contributors, 2016. HIV in the UK - 2016 report.
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/602942/HIV_in_the_UK_report.pdf. (Accessed 31/08/17).
- Lorvick, J., Bluthenthal, R.N., Scott, A., Lou Gilbert, M., Riehm, K.S., Anderson, R.L., Flynn, N.M., Kral, A.H., 2006. Secondary Syringe Exchange Among Users of 23 California Syringe Exchange Programs. *Subst. Use Misuse.* 41, 865-882.
- Marshall, Z., Dechman, M.K., Minichiello, A., Alcock, L., Harris, G.E., 2015. Peering into the literature: A systematic review of the roles of people who inject drugs in harm reduction initiatives. *Drug Alcohol Depend.* 151, 1-14.
- McVeigh, J., Begley, E., 2017. Anabolic steroids in the UK: an increasing issue for public health. *Drugs: Educ. Prev. Policy.* 24, 278-285.
- McVeigh, J., Beynon, C., Bellis, M.A., 2003. New challenges for agency based syringe exchange schemes: analysis of 11 years of data (1991-2001) in Merseyside and Cheshire, United Kingdom. *Int. J. Drug Policy.* 14, 399-405.

McVeigh, J., Chandler, M., Beynon, C., et al., 2007. The injectors that harm reduction forgot. Poster presented at the 18th International Conference on the Reduction of Drug Related Harm, 13-17 May 2007. Warsaw, Poland.

Newland, J., Newman, C., Treloar, C., 2016a. "We get by with a little help from our friends": Small-scale informal and large-scale formal peer distribution networks of sterile injecting equipment in Australia. *International Journal of Drug Policy*. 34, 65-71.

Newland, J., Newman, C., Treloar, C., 2016b. "We get by with a little help from our friends": Small-scale informal and large-scale formal peer distribution networks of sterile injecting equipment in Australia. *Int. J. Drug Policy*. 34, 65-71.

NICE, 2014a. IPED Peer Exchange scheme. <https://www.nice.org.uk/sharedlearning/iped-peer-exchange-scheme>. (Accessed 29/11/2016).

NICE, 2014b. Needle and syringe programmes: NICE public health guidance 52. NICE, London.

Public Health England, Health Protection Scotland, Public Health Wales, Public Health Agency Northern Ireland, 2016. Shooting Up: Infections among people who inject drugs in the UK, 2015. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/567231/Shooting_Up_2016_Update.pdf.

Public Health England, N.I.S., July 2015. Unlinked Anonymous Monitoring Survey of People Who Inject Drugs in contact with specialist services: data tables. People who inject image and performance enhancing drugs. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/537597/UAM_Survey_of_PWID_2016_IPED_data_tables_with_2014_15_data_FINAL.pdf. (Accessed 31/08/17).

Snead, J., Downing, M., Lorvick, J., Garcia, B., Thawley, R., Kegeles, S., Edlin, B.R., 2003.

Secondary syringe exchange among injection drug users. *J. Urban Health*. 80, 330-348.

Tyndall, M.W., Bruneau, J., Brogly, S., Spittal, P., O'Shaughnessy, M.V., Schechter, M.T., 2002.

Satellite needle distribution among injection drug users: Policy and practice in two Canadian cities. *J. Acquir. Immune Defic. Syndr.* 31, 98-105.

Winyard, R., 2005. *Substance Misuse in Primary Care: A Multi-Disciplinary Approach*. Radcliffe Publishing, Oxford, Seattle.

Table 1: Characteristics of study participants by whether they attended a (NSP) and reported collecting equipment for others

		Attended an NSP; n (%)			Among those attending NSP: Collected equipment for others; n (%)		
		Yes	No	<i>p</i>	Yes	No	<i>p</i>
Total number of participants		467	70		157	310	
<i>Demographics</i>							
Age	< 25 years	97 (21%)	20 (29%)	0.334	30 (19%)	67 (22%)	0.275
	25-34 years	203 (44%)	28 (40%)		63 (40%)	140 (45%)	
	>=35 years	167 (36%)	22 (31%)		64 (41%)	103 (33%)	
Gender	Male	457 (98%)	69 (99%)	0.695	153 (97%)	304 (98%)	0.666
Origin of birth	UK born	447 (96%)	65 (93%)	0.289	151 (96%)	296 (95%)	0.726
Ever imprisoned	Yes	116 (25%)	14 (20%)	0.378	45 (29%)	71 (23%)	0.174
<i>Intervention uptake (ever reported)</i>							
Hepatitis B vaccination (at least one dose)	Yes	190 (41%)	23 (33%)	0.212	70 (45%)	120 (39%)	0.222
Tested for hepatitis C	Yes	181 (39%)	20 (29%)	0.101	63 (40%)	118 (38%)	0.666
<i>Infection harms and risks (ever)</i>							
Redness, tenderness and swelling at an injection site	Yes	286 (61%)	24 (34%)	<0.001	107 (68%)	179 (58%)	0.029
Abscess, a sore, or an open wound at an injection site	Yes	74 (16%)	6 (8.6%)	0.111	33 (21%)	41 (13%)	0.029
Anti-HCV	Positive	24 (5.1%)	1 (1.4%)	0.169	12 (7.6%)	12 (3.9%)	0.081
Anti-HIV	Positive	7 (1.5%)	0 (0%)	0.303	3 (1.9%)	4 (1.3%)	0.602
Anti-HBc	Positive	14 (3.0%)	1 (1.4%)	0.457	8 (5.1%)	6 (1.9%)	0.059
<i>Injecting behaviour</i>							
IPED injected in preceding year	Anabolic steroids	389 (94%)	45 (98%)	0.262	130 (93%)	259 (94%)	0.599
	Growth hormone	146 (35%)	14 (30%)	0.521	56 (40%)	90 (33%)	0.142
	Human chorionic gonadotropin	113 (27%)	8 (17%)	0.15	33 (24%)	80 (29%)	0.232
	Melanotan	54 (13%)	3 (6.5%)	0.205	23 (16%)	31 (11%)	0.14
Injected psychoactive substances	Yes, ever	65 (14%)	6 (8.6%)	0.218	30 (19%)	35 (11%)	0.021
	Yes, in preceding year	29 (6.2%)	2 (2.9%)	0.262	17 (11%)	12 (3.9%)	0.011

Ever re-used own needles	Yes	78 (17%)	6 (8.6%)	0.081	33 (21%)	45 (15%)	0.075
Ever used other people's needles	Yes	15 (3.2%)	0 (0%)	0.128	8 (5.1%)	7 (2.3%)	0.1
Ever shared a vial	Yes	56 (12%)	4 (5.7%)	0.12	29 (18%)	27 (9%)	0.002

Table 2: Extent of onward distribution reported by those who collect injecting equipment for others

Number of users who reported collecting for:	n	%	Estimated onward distribution		
			Low ^a	Mid ^b	High ^c
1 other	85	55%	85	85	85
2-9 others	42	27%	84	210	378
10-19 others	7	4.6%	70	105	133
20+ others	20	13%	400	600	1,000
Total	154	100%	639	1,000	1,596

^a calculated using bottom of the category ranges, i.e. 1, 2, 10, & 20

^b calculated using mid-point of the category ranges i.e. 1, 5 & 15 with 30 used for 20+ category

^c calculated using top of the category ranges, i.e. 1, 9 & 19, with 50 is used for the 20+ category