



## LJMU Research Online

**Brandariz-de-Pedro, G, Heyes, DJ, Hardman, SJO, Shanmugam, M, Jones, AR, Weber, S, Nohr, D, Scrutton, NS and Fielding, AJ**

**Direct Evidence of an Excited-State Triplet Species upon Photoactivation of the Chlorophyll Precursor Protochlorophyllide**

<http://researchonline.ljmu.ac.uk/id/eprint/9994/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Brandariz-de-Pedro, G, Heyes, DJ, Hardman, SJO, Shanmugam, M, Jones, AR, Weber, S, Nohr, D, Scrutton, NS and Fielding, AJ (2017) Direct Evidence of an Excited-State Triplet Species upon Photoactivation of the Chlorophyll Precursor Protochlorophyllide. Journal of Physical Chemistry Letters. 8 (6).**

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

# Direct Evidence of an Excited State Triplet Species upon Photoactivation of the Chlorophyll Precursor Protochlorophyllide

*Guillem Brandariz-de-Pedro,<sup>†,‡</sup> Derren J. Heyes,<sup>\*,‡</sup> Samantha J. O. Hardman,<sup>‡</sup> Muralidharan Shanmugam,<sup>‡</sup> Alex R. Jones,<sup>†,‡</sup> Stefan Weber,<sup>§</sup> Daniel Nohr,<sup>§</sup> Nigel S. Scrutton,<sup>‡</sup> Alistair J. Fielding,<sup>\*,†</sup>*

<sup>†</sup>School of Chemistry and the Photon Science Institute, University of Manchester, Oxford Road, Manchester M13 9PL, U.K.

<sup>‡</sup>Manchester Institute of Biotechnology, University of Manchester, 131 Princess Street, Manchester M1 7DN, U.K.

<sup>§</sup>Institute of Physical Chemistry, Albert-Ludwigs-Universität Freiburg, 79104 Freiburg, Germany

## AUTHOR INFORMATION


### Corresponding Author

\*E-mail: [Alistair.Fielding@manchester.ac.uk](mailto:Alistair.Fielding@manchester.ac.uk)

\*E-mail: [Derren.Heyes@manchester.ac.uk](mailto:Derren.Heyes@manchester.ac.uk)


ABSTRACT: The chlorophyll precursor protochlorophyllide (Pchl<sub>id</sub>e), which is the substrate for the light-driven enzyme protochlorophyllide oxidoreductase, has unique excited-state properties that facilitate photocatalysis. Previous time-resolved spectroscopy measurements have implied that a long-lived triplet state is formed during the excited-state relaxation of Pchl<sub>id</sub>e, although direct evidence for its existence is still lacking. Here, we use time-resolved electronic paramagnetic resonance (EPR) in combination with time-resolved absorption measurements at a range of temperatures (10-290 K), solvents, and oxygen concentrations to provide a detailed characterization of the triplet state of Pchl<sub>id</sub>e. The triplet decays in a biphasic, oxygen-dependent manner, whilst the first reported EPR signature of a Pchl<sub>id</sub>e triplet displays both emissive and absorptive features and an antisymmetric spectrum similar to other porphyrin triplet states. This work demonstrates that the Pchl<sub>id</sub>e triplet is accessible to various cryogenic spectroscopic probes over a range of timescales and paves the way for understanding its potential role in catalysis.

## TOC GRAPHICS



**KEYWORDS** electron paramagnetic resonance, protochlorophyllide, triplet state, photochemistry, time-resolved spectroscopy, excited state

Protochlorophyllide (Pchlde) (Figure 1), a crucial intermediate in the biosynthesis of chlorophyll, is the predominant pigment found in seedlings and dark-grown plants.<sup>1-3</sup> As the substrate for the light-driven enzyme protochlorophyllide oxidoreductase (POR) Pchlde has unique excited state properties that allow it to capture sunlight and trigger the subsequent formation of chlorophyll and development of the plant.<sup>3</sup> The reaction catalyzed by POR involves a light-driven hydride transfer from NADPH to the C17 position of Pchlde,<sup>4,5</sup> followed by a thermally-activated proton transfer from a conserved Tyr residue to the C18 position.<sup>6</sup> Catalysis is dependent on excited-state processes in the Pchlde molecule, which occur on an ultrafast, picosecond, timescale. Previous time-resolved absorption and fluorescence spectroscopy measurements on the isolated Pchlde substrate have shown that it is an intrinsically reactive molecule with multi-phasic excited-state dynamics.<sup>3,7-15</sup>




**Figure 1.** Decay lifetimes of protochlorophyllide's excited states.

The excited-state dynamics depend strongly on solvent polarity<sup>7-9</sup> and have been interpreted by using both a branched model<sup>8-13</sup> and a sequential model<sup>7</sup> (Figure 1). However, both models involve the formation of an excited-state intramolecular charge transfer (ICT) state<sup>7-15</sup> and formation of a putative long-lived triplet state by intersystem crossing on a nanosecond timescale.<sup>7,12</sup> The triplet state has absorption bands at approximately 530 and 890 nm and is formed with a significant quantum yield (0.23-0.34).<sup>7,12</sup> However, the existence of this triplet species has only been based on circumstantial evidence, such as the long lifetime and absorption features in the near infra-red region.<sup>12</sup> Moreover, it is unclear whether the triplet state is directly involved in the reaction mechanism of POR<sup>3,6,16</sup> or if the enzyme has a photoprotective role in preventing non-productive side reactions of the highly reactive triplet intermediate.<sup>12</sup> As triplet states are paramagnetic we have used time-resolved electron paramagnetic resonance (EPR) spectroscopy to provide the first reported direct characterization of the excited-state Pchlide triplet species. These measurements have been combined with cryogenic time-resolved absorption measurements to investigate the thermodynamic properties and oxygen-dependence of triplet formation and decay.

Using EPR spectroscopy it is only possible to observe the triplet state at cryogenic temperatures. Hence, initially it was important to provide a direct comparison of the excited-state dynamics of Pchlide at both cryogenic and room temperatures by using time-resolved absorption measurements after laser excitation at 450 nm (ps to  $\mu$ s timescale) as described in the Supporting Information. The time-resolved difference spectra from these measurements (Figures S1-S8) were fitted to a sequential kinetic model by global analysis to yield evolution associated difference spectra (EADS) (Figure 2).<sup>16</sup> The time-resolved spectral changes at a range of temperatures between 10 K and room temperature, and in the presence and

absence of oxygen could all be fitted to four components with only minor differences in the lifetimes of the various excited state species (Table 1). The first three excited-state components have been described previously and correspond to the formation of an ICT state from the  $S_1$  excited state ( $\sim 5$  ps), solvation of the ICT state ( $\sim 200$  ps) and decay of the  $S_1/S_{ICT}$  excited state into the long-lived triplet state on the nanosecond timescale ( $\sim 4$  ns).<sup>7-13</sup> Hence, it appears that temperature and presence of oxygen only have a minimal effect on these three excited-state processes, which are directly related to the formation of the triplet state. The remaining long-lived component ( $\tau_4$ ), which relates to the decay of the triplet state, occurs on longer timescales than the  $80 \mu\text{s}$  time window of the pump-probe measurements.



**Figure 2.** Evolution associated difference spectra (EADS) resulting from a global analysis of the time-resolved absorption data for an aqueous, anaerobic Pchlide sample after excitation at 450 nm at 10 K. The EADS were obtained from fitting the time-resolved absorption data to a sequential exponential model as described in the supplementary information. The triplet decay lifetime was set to infinity on the  $80 \mu\text{s}$  timeframe of the measurement.

**Table 1.** Lifetimes of the  $S_1$  ( $\tau_1$ ),  $S_{ICT}$  ( $\tau_2$ ), solvated  $S_{ICT}$  ( $\tau_3$ ) and triplet excited states of Pchlride ( $\tau_4$  and  $\tau_5$ ) measured in both aerobic and anaerobic aqueous buffer solutions at 10, 80, 150 and 290 K. For experimental conditions see Supporting Information.


	T (K)	$\tau_1$ (ps)	$\tau_2$ (ps)	$\tau_3$ (ns)	$\tau_4$ (ms)	$\tau_5$ (ms)
anaerobic	10	$4.0 \pm 0.2$	$215 \pm 5$	$5.8 \pm 0.1$	$6.9 \pm 0.1$	$24.9 \pm 0.2$
	80	$6.3 \pm 0.3$	$276 \pm 5$	$4.9 \pm 0.1$	$7.5 \pm 0.1$	$22.3 \pm 0.2$
	150	$5.0 \pm 0.2$	$170 \pm 3$	$4.1 \pm 0.1$	$6.4 \pm 0.1$	$19.4 \pm 0.1$
	290	$3.4 \pm 0.4$	$180 \pm 6$	$3.5 \pm 0.1$	$0.35 \pm 0.01$	$1.82 \pm 0.01$
aerobic	10	$4.6 \pm 0.2$	$205 \pm 8$	$5.9 \pm 0.1$	$3.1 \pm 0.2$	-
	80	$4.7 \pm 0.2$	$597 \pm 20$	$6.0 \pm 0.1$	$3.6 \pm 0.1$	-
	150	$5.6 \pm 0.2$	$278 \pm 9$	$5.5 \pm 0.1$	$3.7 \pm 0.2$	-
	290	$3.8 \pm 0.2$	$364 \pm 26$	$4.9 \pm 0.2$	$0.11 \pm 0.01$	-

In order to investigate the kinetics of triplet decay in more detail the time-resolved absorption measurements were extended to the ms timescale by using laser flash photolysis experiments (Figures S9-S16). Consequently, the triplet state of Pchlride was found to decay in a biphasic process with  $\tau_4$  values of approximately 7 ms and  $\tau_5$  values of approximately 22 ms under cryogenic and anaerobic conditions (Figure 3A, Table 1). However, in the presence of oxygen the triplet decay kinetics become faster and more monoexponential in nature, displaying  $\tau_4$  values of  $\sim 3.5$  ms at cryogenic temperatures. The oxygen dependence was investigated further by following the recovery of the Pchlride ground state (absorbance at 630 nm) over a range of oxygen concentrations at room temperature (Figure 3B). The triplet decay kinetics show a strong dependence on oxygen concentration, confirming that the triplet excited-state of Pchlride can interact with molecular oxygen, presumably to yield singlet oxygen or other reactive oxygen species

via triplet-triplet energy transfer.<sup>17,18</sup> In addition, the observed lifetimes show no dependence on temperature at cryogenic temperatures (Figure S33) suggesting that the decay of the triplet state is an adiabatic process in frozen samples.

As previous studies have shown that the excited-state dynamics of Pchl<sub>id</sub>e depend strongly on solvent polarity<sup>7-9</sup> the transient absorption measurements were repeated in alcohol solutions. The ultrafast steps leading to the formation of the triplet state are very similar to those in water and show no noticeable dependence on oxygen or temperature (Figure S34, Table S1). However, the decay of the Pchl<sub>id</sub>e triplet state now appears to follow single, monoexponential kinetics under both aerobic and anaerobic conditions with a lifetime of ~3 ms at cryogenic temperatures. Moreover, at higher temperatures the triplet excited state decays much more rapidly in polar solvent (lifetime of ~2 μs) than in water (lifetime of ~0.1 ms). It has previously been shown that dimers/aggregates of Pchl<sub>id</sub>e, formed in water-based buffers, display different excited state dynamics to monosolvated Pchl<sub>id</sub>e found in polar solvents.<sup>19,20</sup> Hence, it is possible that the differences in aggregation state of Pchl<sub>id</sub>e may also explain the changes in triplet decay kinetics observed in the present work.





**Figure 3.** (A) EADS resulting from a global analysis of the laser flash photolysis data for an aqueous, anaerobic Pchl<sub>a</sub> sample after excitation at 450 nm at 10 K. The EADS were obtained from fitting the data to a sequential exponential model as described in the supplementary information. (B) Lifetime values ( $\tau_4$  and  $\tau_5$ ) obtained from fitting a double exponential decay to ground state recovery (630 nm) flash photolysis time traces. Samples consisted of Pchl<sub>a</sub> dissolved in mixtures of anaerobic and aerobic aqueous buffer at room temperature.


In order to provide direct experimental evidence of the existence of an excited-state triplet of Pchl<sub>id</sub> time-resolved EPR has been used after photoexcitation with laser pulses at 532 nm (Figure 4) and 450 nm (Figure S36). The EPR signature of the photoexcited triplet state Pchl<sub>id</sub> was obtained from integrating direct-detection EPR time traces over a range of magnetic field positions. The EPR spectrum displays both emissive and absorptive features and an antisymmetric lineshape (Figure 4A), which are all features that are characteristic of a triplet state.<sup>21-23</sup> The lifting of the degeneracy of spin states in the absence of an applied magnetic field is referred to as zero-field splitting (ZFS). The value of the ZFS parameter is usually defined via D and E parameters. D describes the axial component of the magnetic dipole-dipole interaction, and E the transversal component. The values of the ZFS splitting parameters obtained from a simulation of the spectrum (Figure S37) are of a rhombic symmetry and were found to be  $|D| = 860 \pm 8$  and  $|E| = 182 \pm 5$  MHz. These values are comparable to those of chlorophyll *a* ( $|D| = 818 \pm 24$  and  $|E| = 120 \pm 6$  MHz) and chlorophyll *b* ( $|D| = 878 \pm 27$  and  $|E| = 156 \pm 3$  MHz)<sup>24</sup>, suggesting that the energy states of the triplet state of Pchl<sub>id</sub> are similar to those found in the chlorophylls. Changes in the E value reflect some differences in asymmetry. Density functional theory calculations incorrectly predicted ZFS parameters as  $|D| = 648$  and  $|E| = 166$  MHz, although this error might be due to the inability of the method to accurately take into account various molecular orbitals in highly delocalized systems, such as photoexcited Pchl<sub>id</sub>.<sup>25</sup>

Electron spin echo (ESE) field-swept spectra (Figure 4A) measured at different delay-after-flash (DAF) values reflect the anisotropic decay of the triplet sub-levels of the Pchl<sub>id</sub> triplet state, as previously observed in other organic triplets.<sup>26-28</sup> Triplet sub-level

decay rates were calculated from measuring the intensity of the two-pulse spin echo EPR signal at increasing DAF values. A representation of the ESE intensity vs the DAF is shown in Figure 4B. A double exponential decay with a positive and a negative component was fit to the time evolution traces. These were measured at the high-field  $Z^+$  orientation of the triplet state EPR spectrum for Pchl $a$  samples in aerobic and anaerobic aqueous and alcohol solutions, providing the decay constants from  $T_0$  for that orientation as per the following formula:

$$I = A_1 \exp(-t/\tau_1) + A_0 \exp(-t/\tau_0)$$

with  $I$  being the ESE intensity,  $t$  the DAF,  $|A_1|$  and  $|A_0|$  being the initial relative populations of  $T_{+1}/T_{-1}$  and  $T_0$ , with  $\tau_1$  and  $\tau_0$  as their respective decay lifetimes.  $A_0$  is positive for the “+” canonical orientations and negative for the “-” orientations, representing absorption and emission from  $T_0$  respectively. The opposite is true for  $A_1$  and  $T_{+1}/T_{-1}$ . The triplet EPR decay lifetimes in different solvents show the same trends as the optical experiments, with faster decays in the alcohol solutions and in the presence of oxygen. However, it should be noted that the EPR signal intensity for the aerobic aqueous sample was too weak to measure accurately.



**Figure 4.** (A) ESE field-swept spectra of Pchlride recorded at 10 K in anaerobic aqueous buffer at different DAF values displaying an anisotropic decay of the triplet state spectrum features. (B) ESE-detected kinetics of the Pchlride triplet in anaerobic aqueous buffer and both anaerobic and aerobic alcohol solutions at 10 K at the high-field  $Z^+$  canonical orientation. Lifetimes correspond to the decay from  $T_0$ . **Inset:** Pulse sequence for the EPR two-pulse electron spin echo (ESE) experiment consisting of a laser pulse (in green), microwave pulses (in blue at a delay after flash DAF and separated by a delay  $\tau$ ) and the generated spin echo signal.

This work provides the first direct evidence that a long-lived triplet state is formed during the excited-state relaxation of Pchlride. Time-resolved EPR measurements confirm that the Pchlride

triplet, which forms on the nanosecond timescale and decays on the millisecond timescale, is similar to other porphyrin triplet states. The Pchl<sub>ide</sub> excited-state triplet species has been shown to react with oxygen, suggesting that highly reactive oxygen species could be formed, which may lead to photo-damage in the cell. Hence, it is likely that specific chromophore-protein interactions occur when Pchl<sub>ide</sub> is bound to the enzyme, which would help to minimize any undesirable effects of Pchl<sub>ide</sub> triplet formation. These findings open up an exciting new range of pulsed EPR spectroscopy studies such as ESE envelope modulation (ESEEM), hyperfine sublevel correlation (HYSCORE) or electron nuclear double resonance (ENDOR), which will be essential to gain further information on the electronic couplings of Pchl<sub>ide</sub> in the excited triplet state. In addition, measurements on POR-bound Pchl<sub>ide</sub> will provide important insights into how the enzyme active site affects such electronic properties and the potential role of the triplet state in POR catalysis.

## ASSOCIATED CONTENT

**Supporting Information.** The following files are available free of charge. Detailed experimental methods, raw transient absorption data, thermodynamic data, EPR data (PDF).

## AUTHOR INFORMATION

### Notes

The authors declare no competing financial interests.

## ACKNOWLEDGMENTS

G.B.-P. acknowledges Early Stage Researcher funding from the European Union's Seventh Framework Programme FP7-PEOPLE-2013-ITN through the 'MAGnetic Innovation in Catalysis' (MAGIC) Initial Training Network (Grant agreement no. 606831). We would like to acknowledge the use of the EPSRC EPR National Service in carrying out this work. NSS is an Engineering and Physical Sciences Research Council (EPSRC) Established Career Fellow (EP/J020192/1). Ultrafast spectroscopy was performed at the Ultrafast Biophysics Facility, Manchester Institute of Biotechnology, as funded by BBSRC Alert14 Award BB/M011658/1.

## REFERENCES

- (1) Heyes, D. J.; Hunter, C. N. Making Light Work of Enzyme Catalysis: Protochlorophyllide Oxidoreductase. *Trends Biochem. Sci.* **2005**, *30* (11), 642–649.
- (2) Masuda, T.; Takamiya, K. I. Novel Insights into the Enzymology, Regulation and Physiological Functions of Light-Dependent Protochlorophyllide Oxidoreductase in Angiosperms. *Photosynthesis Research*. 2004, pp 1–29.
- (3) Scrutton, N. S.; Groot, M. L.; Heyes, D. J. Excited State Dynamics and Catalytic Mechanism of the Light-Driven Enzyme Protochlorophyllide Oxidoreductase. *Phys. Chem. Chem. Phys.* **2012**, *14* (25), 8818–8824.
- (4) Heyes, D. J.; Heathcote, P.; Rigby, S. E. J.; Palacios, M. a; van Grondelle, R.; Hunter, C. N. The First Catalytic Step of the Light-Driven Enzyme Protochlorophyllide Oxidoreductase Proceeds via a Charge Transfer Complex. *J. Biol. Chem.* **2006**, *281* (37), 26847–26853.

- (5) Heyes, D. J.; Sakuma, M.; de Visser, S. P.; Scrutton, N. S. Nuclear Quantum Tunneling in the Light-Activated Enzyme Protochlorophyllide Oxidoreductase. *J. Biol. Chem.* **2009**, *284* (6), 3762–3767.
- (6) Menon, B. R. K.; Waltho, J. P.; Scrutton, N. S.; Heyes, D. J. Cryogenic and Laser Photoexcitation Studies Identify Multiple Roles for Active Site Residues in the Light-Driven Enzyme Protochlorophyllide Oxidoreductase. *J. Biol. Chem.* **2009**, *284* (27), 18160–18166.
- (7) Sytina, O. A.; van Stokkum, I. H. M.; Heyes, D. J.; Hunter, C. N.; van Grondelle, R.; Groot, M. L. Protochlorophyllide Excited-State Dynamics in Organic Solvents Studied by Time-Resolved Visible and Mid-Infrared Spectroscopy. *J. Phys. Chem. B* **2010**, *114* (C), 4335–4344.
- (8) Dietzek, B.; Maksimenka, R.; Siebert, T.; Birckner, E.; Kiefer, W.; Popp, J.; Hermann, G.; Schmitt, M. Excited-State Processes in Protochlorophyllide a – a Femtosecond Time-Resolved Absorption Study. *Chem. Phys. Lett.* **2004**, *397* (1–3), 110–115.
- (9) Dietzek, B.; Kiefer, W.; Hermann, G.; Popp, J.; Schmitt, M. Solvent Effects on the Excited-State Processes of Protochlorophyllide: A Femtosecond Time-Resolved Absorption Study. *J. Phys. Chem. B* **2006**, *110* (9), 4399–4406.
- (10) Dietzek, B.; Kiefer, W.; Yartsev, A.; Sundström, V.; Schellenberg, P.; Grigaravicius, P.; Hermann, G.; Popp, J.; Schmitt, M. The Excited-State Chemistry of Protochlorophyllide a: A Time-Resolved Fluorescence Study. *ChemPhysChem* **2006**, *7* (8), 1727–1733.
- (11) Schmitt, M.; Dietzek, B.; Hermann, G.; Popp, J. Femtosecond Time-Resolved

- Spectroscopy on Biological Photoreceptor Chromophores. *Laser Photonics Rev.* **2007**, *1* (1), 57–78.
- (12) Dietzek, B.; Tschierlei, S.; Hermann, G.; Yartsev, A.; Pascher, T.; Sundström, V.; Schmitt, M.; Popp, J. Protochlorophyllide a: A Comprehensive Photophysical Picture. *Chemphyschem* **2009**, *10* (1), 144–150.
- (13) Dietzek, B.; Tschierlei, S.; Hanf, R.; Seidel, S.; Yartsev, A.; Schmitt, M.; Hermann, G.; Popp, J. Dynamics of Charge Separation in the Excited-State Chemistry of Protochlorophyllide. *Chem. Phys. Lett.* **2010**, *492* (1–3), 157–163.
- (14) Colindres-Rojas, M.; Wolf, M. M. N.; Gross, R.; Seidel, S.; Dietzek, B.; Schmitt, M.; Popp, J.; Hermann, G.; Diller, R. Excited-State Dynamics of Protochlorophyllide Revealed by Subpicosecond Infrared Spectroscopy. *Biophys. J.* **2011**, *100* (1), 260–267.
- (15) Hanf, R.; Fey, S.; Dietzek, B.; Schmitt, M.; Reinbothe, C.; Reinbothe, S.; Hermann, G.; Popp, J. Protein-Induced Excited-State Dynamics of Protochlorophyllide. *J. Phys. Chem. A* **2011**, *115*, 7873–7881.
- (16) Heyes, D. J.; Hardman, S. J. O.; Hedison, T. M.; Hoeven, R.; Greetham, G. M.; Towrie, M.; Scrutton, N. S. Excited-State Charge Separation in the Photochemical Mechanism of the Light-Driven Enzyme Protochlorophyllide Oxidoreductase \*\*. *Angew. Chemie - Int. Ed.* **2015**, *54* (5), 1512–1515.
- (17) Patterson, L. K.; Porter, G.; Topp, M. R. Oxygen Quenching of Singlet and Triplet States. *Chem. Phys. Lett.* **1970**, *7* (6), 612–614.



- (18) Gijzeman, O. L. J.; Kaufman, F.; Porter, G. Oxygen Quenching of Aromatic Triplet States in Solution. Part 1. *J. Chem. Soc. Faraday Trans. 2 Mol. Chem. Phys.* **1973**, *69* (0), 708–720.
- (19) Lebedev, N. N.; Krasnovsky, A. A.; Litvin, F. F. Phosphorescence of Protochlorophyll(ide) and Chlorophyll(ide) in Etiolated and Greening Bean Leaves - Assignment of Spectral Bands. *Photosynth. Res.* **1991**, *30* (1), 7–14.
- (20) Heyes, D. J.; Hardman, S. J. O.; Mansell, D.; Gardiner, J. M.; Scrutton, N. S. Mechanistic Reappraisal of Early Stage Photochemistry in the Light-Driven Enzyme Protochlorophyllide Oxidoreductase. *PLoS One* **2012**, *7* (9), e45642.
- (21) Hutchison, C. A.; Mangum, B. W. Paramagnetic Resonance Absorption in Naphthalene in Its Phosphorescent State. *J. Chem. Phys.* **1958**, *29* (1), 952–953.
- (22) Yager, W. A.; Wasserman, E.; Cramer, R. M. R. ESR Observation of  $\Delta m=1$  Transitions of Triplet States in Glasses. **1962**, *37* (1), 1148–1149.
- (23) Waals, J. H. EPR of Photo-Excited Triplet States: A Personal Account. *Applied Magnetic Resonance*. 2001, pp 545–561.
- (24) Thurnauer, M. C.; Norris, J. R. The Ordering of the Zero Field Triplet Spin Sublevels in the Chlorophylls. A Magnetophotoselection Study. *Chem. Phys. Lett.* **1977**, *47* (1), 100–105.
- (25) Cohen, A. J.; Mori-Sánchez, P.; Yang, W. Challenges for Density Functional Theory. *Chem. Rev.* **2012**, *112* (1), 289–320.

- (26) Schmidt, J.; Antheunis, D. A.; van der Waals, J. H. The Dynamics of Populating and Depopulating the Phosphorescent Triplet State as Studied by Microwave Induced Delayed Phosphorescence. *Mol. Phys.* **1971**, *22* (1), 1–17.
- (27) Lin, T. S. Electron Spin Echo Spectroscopy of Organic Triplets. *Chem. Rev* **1984**, *84* (1), 1–15.
- (28) Di Valentin, M.; Ceola, S.; Agostini, G.; Giacometti, G. M.; Angerhofer, A.; Crescenzi, O.; Barone, V.; Carbonera, D. Pulse ENDOR and Density Functional Theory on the Peridinin Triplet State Involved in the Photo-Protective Mechanism in the Peridinin-Chlorophyll a-Protein from *Amphidinium Carterae*. *Biochim. Biophys. Acta - Bioenerg.* **2008**, *1777* (3), 295–307.